

Human McI-1 Alexa Fluor® 700-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 542808 Catalog Number: FAB828N

100 µg

DESCRIPTION		
Species Reactivity	Human	
Specificity	Detects human McI-1.	
Source	Monoclonal Mouse IgG _{2B} Clone # 542808	
Purification	Protein A or G purified from hybridoma culture supernatant	
Immunogen	E. coli-derived recombinant human Mcl-1 Met1-Gly230 Accession # Q07820	
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm	
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS			
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.			
Knockout Validated	Optimal dilution of this antibody should be experimentally determined.		
Western Blot	Optimal dilution of this antibody should be experimentally determined.		
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.		

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied	

BACKGROUND

Mcl-1 (myeloid cell leukemia-1; also known as Bcl-2-like protein 3) is a member of the Bcl-2 family of proteins. Alternative splicing creates two distinct isoforms: 40 kDa Mcl-1L (long; 350 amino acids (aa)) enhances cell survival by inhibiting apoptosis, while 32 kDa Mcl-1S (short; 271 aa with divergence in the last 41 aa) promotes apoptosis. The elimination of Mcl-1L is a required step for DNA damage-induced apoptosis. Mcl-1 can be modified by phosphorylation on S121 and T163 by JNK, which triggers apoptosis, or polyubiquitination, which enhances degradation of Mcl-1. Within the first 230 aa, human Mcl-1 shares ~68% aa identity with mouse and rat Mcl-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/23/2025 Page 1 of 1

China | info.cn@bio-techne.com TEL: 400.821.3475