

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human GDF-5 in direct ELISA.
Source	Monoclonal Mouse IgG _{2B} Clone # 1065032
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	E. coli-derived recombinant human GDF-5 Ala382-Arg501 Accession # P43026
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Growth Differentiation Factor 5 (GDF-5), also known as cartilage-derived morphogenetic protein 1 (CDMP-1) and BMP-14, is a member of the bone morphogenetic protein (BMP) family which belongs to the transforming growth factor β (TGF- β) superfamily. GDF-5 is synthesized as a large precursor protein that consists of an N-terminal 19 amino acid (aa) signal sequence, a 362 aa pro region and a 120 aa C-terminal mature peptide. Mature GDF-5 is a homodimeric protein which contains the characteristic seven conserved cysteine residues. GDF-5, GDF-6 and GDF-7, which share 80-86% identity, define a subgroup within the BMP family. Like other TGF- β superfamily proteins, GDF-5 is highly conserved across species. At the amino acid sequence level, mature human and mouse GDF-5 are 98% identical. It has been reported that GDF-5 has multiple functions including regulation of myogenesis, regulation of chondrogenesis, bone morphogenesis, and neuron differentiation and survival. GDF-5 response is mediated by the formation of hetero-oligomeric complexes of type I (BMPRI-B) and type II (BMPRII or Activin R-II) serine/threonine kinase receptors, and the activation of Smad proteins (Smad 1, 5, and 8).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.