

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human FAM20B in direct ELISAs.
Source	Recombinant Monoclonal Mouse IgG ₁ Clone # 1018512
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human FAM20B Asn27-Leu409 Accession # O75063
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

FAM20B is a member of the FAM20 protein family that has three members in mammals (FAM20A, FAM20B, FAM20C) (1). Human FAM20B shares 97% amino acid sequence identity with mouse and rat FAM20B. FAM20B localizes to the Golgi apparatus where it phosphorylates the xylose residue in the glycosaminoglycan (GAG)-protein linkage region of proteoglycans, which leads to enhanced GAG biosynthesis (2). Accordingly, chondroitin sulfate and heparan sulfate production is increased when FAM20B is over-expressed and reduced following FAM20B knockdown (2). FAM20B knockout mice display embryonic lethality at day E13.5, suggesting that FAM20B has an important role during development (3). Furthermore, in zebrafish, FAM20B mutations result in reduced cartilage matrix production and skeletal defects (4). The enzymatic activity of recombinant human FAM20B is measured using a phosphatase-coupled method (5).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.