

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Neuropeptide Y/NPY in direct ELISAs.
Source	Monoclonal Mouse IgG _{2A} Clone # 904032
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Neuropeptide Y/NPY conjugated to KLH YPSKPDNPGEDAPAEDMARYYSALRHYINLITRQRY Accession # P01303
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Neuropeptide Y (NPY) is a 36 amino acid peptide that was isolated from hypothalamus in porcine brain in 1982 and lately it belongs to a family of peptides which include Pancreatic Polypeptide (PP) and Peptide YY (PYY) which exert their pharmacological action via interaction with G-protein coupled receptors Y1, Y2, Y4, Y5 and y6. NPY is the most abundant peptide in brain and in nervous system NPY functions as a neurotransmitter regulating many processes including memory and learning, pain, fat storage and blood pressure. NPY also regulates stress by stimulating secretion of corticotropin-releasing hormone in brain. It appears there is a correlation between the increased levels of NPY gene expression in hippocampus and epileptic seizures. Cocaine reduces the levels of NPY and such a decrease is thought to be related to depression and anxiety. NPY receptors are rhodopsin-like G-protein coupled receptors (GPCR) coupled to G_i or G_o proteins, which inhibit adenylate cyclase and reduce cAMP accumulation and modulate Calcium and Potassium channels.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.