

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Reelin in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 940209
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Reelin Ser1221-Gln2666 Accession # P78509
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Reelin is a secreted modular glycoprotein that exhibits serine protease activity and is crucial for brain development and function (1-3). It is composed of an N-terminal Reelin domain, eight EGF-like Reelin repeats (RR), and a highly basic C-terminal region (4-6). The N-terminal fragment is suggested to mediate dimerization/oligomerization and receptor recognition, the midpiece receptor binding, and the C-terminal fragment receptor signaling and recognition (1, 5, 7-9). Human Reelin is synthesized as a 3460 amino acid (aa) precursor protein with a molecular weight of approximately 410 kDa (4). During processing, it can be cleaved between RR2 and RR3 or between RR6 and RR7, producing a 180 kDa and a 330 kDa peptide, respectively (1, 6, 10). Within shared regions in the central fragment, human Reelin shares 95% aa sequence identity with mouse and rat Reelin.

Reelin is secreted by Cajal-Retzius cells in the embryo (1, 4, 11). In the adult, it is expressed in the subventricular zone, rostral migratory stream, olfactory bulb, and in the CA1, CA3, and dentate gyrus regions of the hippocampus, as well as in cerebellar granule cells, pyramidal cells of the entorhinal cortex, GABA interneurons, and glial cells (1, 6, 12, 13). Reelin utilizes the receptors VLDLR and ApoE R2, which have been suggested to have divergent roles in Reelin-mediated neuronal migration (1, 2, 6, 12). It has also been shown to interact with Integrin $\alpha 3 \beta 1$ and APP (1, 6, 14, 15). During cortical plate development, Reelin controls cell-cell interactions critical for proper neuronal migration and positioning (1, 2, 4, 5, 11, 12, 16). In the adult, it plays a role in dendrite growth and maturation, and synapse formation (2, 6, 15). Additionally, Reelin has been shown to modulate synaptic transmission and plasticity by regulating the subunit composition and conductivity of NMDA receptors (2, 6, 17). Mutation of the *RELN* gene results in lissencephaly with cerebellar hypoplasia (11, 18). In addition, abnormal Reelin expression in the brain has been associated with a variety of cognitive pathological conditions including autism, schizophrenia, bipolar disorder, major depression, and Alzheimer's disease (1, 6, 11, 13, 19, 20). Peripherally, Reelin is important in the development of neuromuscular junctions. But instead of utilizing the locally expressed ApoE R2 and VLDLR, this function requires the serine protease activity of Reelin (3, 21).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.