


DESCRIPTION	
Species Reactivity	Human
Specificity	Stains human APJ transfectants in flow cytometry but not the parental cell line.
Source	Monoclonal Mouse IgG ₃ Clone # 72133
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	NS0 mouse myeloma cell line transfected with APJ Met1-Asp380 Accession # P35414
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

APJ, also known as Apelin Receptor, GPCR-34 and AGTRL-1, is a G-protein coupled receptor that is related to the angiotensin receptor. It is primarily expressed in coronary artery endothelium and binds the endogenous ligand, Apelin. APJ can also function as a co-receptor for HIV.