

DESCRIPTION


Species Reactivity	Human
Specificity	Detects Integrin α 1/CD49a by direct elisa.
Source	Recombinant Monoclonal Rabbit IgG Clone # 1090A
Purification	Protein A or G purified from cell culture supernatant
Immunogen	CHO-S derived rhITGA-1(1-1141/GSL/aT)/C-His + rhITGB-1(21-728/GSL/bT) Accession # P56199
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Integrin α 1 (also VLA-1, CD49a and Laminin and Collagen Receptor) is a 190-210 kDa member of the integrin alpha chain family of molecules. It is found on smooth muscle cells, osteoblasts, adipocytes and intestinal epithelium. Integrin α 1 forms a noncovalent heterodimer with Integrin β 1, and serves as a divalent-cation dependent receptor for collagen types I, IV, VI, XIII and XVI. It also binds the diarrhea-associated NSP4 enterotoxin of rotavirus. Mature human Integrin α 1 is a 1151 amino acid (aa) type I transmembrane glycoprotein that contains a 1113 aa extracellular domain (ECD) and a 15 aa cytoplasmic tail. The ECD contains one vWFA/I-domain (aa 147-360) that binds collagen, plus multiple divalent cation binding sites. Potential splice variants exist that show a two and four aa substitution for aa 765-1179. Over aa 29-1141 (the ECD), human Integrin α 1 shares 88% aa identity with mouse Integrin α 1.