

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Dihydrolipoamide Dehydrogenase/DLD in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 1016619
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived human Dihydrolipoamide Dehydrogenase/DLD Ala36-Phe509 Accession # P09622
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Dihydrolipoamide dehydrogenase (DLD), also known as LADH, is an NAD-dependent oxidoreductase in the mitochondrial matrix (1). DLD serves as the E3 subunit of four mitochondrial enzyme complexes: pyruvate dehydrogenase, alpha-ketoglutarate dehydrogenase, branched chain alpha-ketotacid dehydrogenase, and the glycine cleavage system (2, 3). It is active as a 120 kDa dimer that catalyzes oxidation within these enzyme complexes. Several mutations of human DLD have been described, some of which contribute to the loss of respiratory function during oxidative stress (4, 5). DLD mutations located at the interface between dimer subunits can impair dimer formation (5, 6). The involvement of DLD in the regulation of lipid peroxidation and lactate metabolism is important for mouse hippocampal neuroblast proliferation and hamster sperm capacitation, respectively (7, 8). DLD polymorphisms in insects can increase their resistance to the pesticide phosphine gas, while they can increase the sensitivity to arsenite in the nematode *C. elegans* (9). Mature human DLD shares 95-96% amino acid (aa) sequence identity with hamster, mouse, and rat DLD. Alternative splicing generates additional human DLD isoforms that lack the N-terminal 99 aa or carry a deletion of aa 147-194.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.