

Mouse Caspase-11/Caspase-4 Alexa Fluor® 750-conjugated Antibody

Monoclonal Rabbit IgG Clone # 1172C Catalog Number: FAB8648S

ımber: FAB8648S 100 µg

DESCRIPTION	
Species Reactivity	Mouse
Specificity	Detects mouse Caspase-11/Caspase-4 in Western blots.
Source	Monoclonal Rabbit IgG Clone # 1172C
Purification	Protein A or G purified from cell culture supernatant
Immunogen	HEK293 human embryonic kidney cell line transfected with mouse Caspase-11/Caspase-4 Accession # P70343
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

China | info.cn@bio-techne.com TEL: 400.821.3475

PREPARATION AND STORAGE	
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUNE

Mouse Caspase-11 (commonly referred as Caspase-4) is a member of the cysteine-aspartic acid protease family known as caspases that function in initiating apoptosis or activating proinflammitory cytokines. Mouse Caspase-11 is expressed as a 343 amino acid latent zymogen. The active caspase-11 is cleaved into two subunits, p20 (aa 81-266) and p10 (aa 286-373). Caspase-11 is one of several caspases associated with the inflammasome and plays an important role in the activation of proinflammatory cytokines. Caspase-11 is activated by an autocatalytic mechanism or by cleavage by Caspase-8 in response to ER stress or bacterial infection, and its expression and activation can be induced by exposure to LPS. The active enzyme can cleave and activate Caspase-1 which in turn activates IL-1 beta. Mouse Caspase-11 has a 59 and 89% homology to human and rat Caspase-4, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/23/2025 Page 1 of 1