

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-10 R β in direct ELISAs and Western blots. In direct ELISAs and Western blots, ~10% cross-reactivity with recombinant human (rh) IL-10 R and rhIFN- α/β R1 is observed and no cross-reactivity with rhIFN- γ R1 and rhIFN- γ R2
Source	Monoclonal Mouse IgG ₁ Clone # 90227
Purification	Protein A or G purified from ascites
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-10 R β Met20-Ser220 Accession # Q08334
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-10 R β , also known as IL-10 R2, mediates its biological activities *via* binding to a heteromeric receptor complex consisting of two distinct type II cytokine receptor subunits, the ligand binding IL-10 R α and the IL-10 R β which does not bind IL-10 by itself but is required for signal transduction. The cDNA for human IL-10 R β encodes a 325 amino acid (aa) residue type I transmembrane precursor protein with a 20 aa signal sequence, a 200 aa extracellular region, a 29 aa transmembrane segment and a short 76 aa cytoplasmic domain. Within the extracellular region, there are two 100 aa subdomains that resemble the constant region of immunoglobulins. This structural motif is responsible for the alternative designation of IL-10 R β as CRF2-4 (the 4th member of the cytokine receptor family class II/2). Human and mouse IL-10 R β share approximately 69% aa sequence identity. Binding of the non-covalent IL-10 dimer to two IL-10 R α chains recruits two IL-10 R β chains resulting in the activation and phosphorylation of the signaling cascade involving JAK1, TYK2, and STAT3. IL-10 R β is expressed ubiquitously. IL-10 R β is also a component of the IL-22 receptor complex consisting of the IL-10 R β chain and IL-22 R, another type II cytokine receptor family member.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.