

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human AICL/CLEC-2B in direct ELISA.
Source	Monoclonal Mouse IgG ₁ Clone # 1064935
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese Hamster Ovary cell line, CHO-derived human AICL/CLEC-2B Lys26-His149 Accession # Q92478
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Flow Cytometry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Activation-induced C-type lectin (AICL), also known as CLEC2B, is a 30-35 kDa variably glycosylated type-2 transmembrane member of the C-type lectin-like receptor (CTLR) family. AICL belongs to the subgroup of CLEC2 proteins that also includes CLEC2A/KACL, CLEC2D/LLT, and CD69/CLEC2C, all of which are encoded by the natural killer gene complex (1). Human AICL contains a single C-type lectin domain in its extracellular region and a 7 amino acid cytoplasmic tail (2). AICL is expressed on monocytes, macrophages, and granulocytes (3), and it is upregulated on TLR-activated monocytes and IL-12 + IL-18 activated NK cells (3, 4). AICL is an activating receptor that triggers TNF production by monocytes (3). It binds to NKp80 on NK cells, resulting in NK cell mediated lysis of the AICL expressing monocyte (3). In addition, the AICL-NKp80 axis mediates interactions between activated and resting NK cells (4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.