

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Dopamine D2 R/DRD2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 960710
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human DRD2 synthetic peptide Accession # P14416
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Dopamine receptor D2 (DRD2) is localized to human striatum, motor cortex and neocortex. It is a highly conserved seven transmembrane receptor member of the G-protein coupled receptor 1 family, with three known isoforms. In the striatum, DRD2 suppresses voluntary activity in the striatopallidal pathway, and polymorphisms in this gene are associated with alcohol addiction, smoking behavior, schizophrenia, food addiction and post-traumatic stress disorder as well as myoclonus dystonia. DRD2 has been shown to heterodimerize with DDR4 and 5-HT2A receptors, and traffics between the plasma membrane and intracellular pools and this localization can be modulated by several drugs. DRD2 is localized to human striatum, motor cortex and neocortex.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.