

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human HIPK2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 493918
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human HIPK2 Met765-Ser924 Accession # Q9H2X6
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Homeodomain-interacting protein kinase-2 (HIPK2) is a serine/threonine kinase which interacts with transcription factors, including p53, CREB1, SMAD1, SMAD2, RUNX1 and others, and can act as a corepressor or coactivator depending on the transcription factor and subcellular localization. Post-translational modifications control HIPK2 activity, including phosphorylation, small ubiquitin-like modifier modification, acetylation, and ubiquitination. HIPK2 regulation of cell proliferation makes it a key effector of proliferative diseases including cancer and kidney fibrosis. For example, HIPK2 can repress the proliferative Wnt-1/b-catenin pathway and HIF-1, promote apoptosis by activation of p53 and repression of ΔNp63α, MDM2 and CtBP, and its activity is enhanced by anti-cancer drugs such as cisplatin and etoposide. HIPK2 oncorepression can be blocked by cytoplasmic localization, hypoxia, gene mutation and loss of heterozygosity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.