

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ADAM28 in direct ELISA.
Source	Monoclonal Rabbit IgG ₁ Clone # 2248B
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Chinese Hamster Ovary cell line, CHO-derived human ADAM28 protein Leu72 & Leu191-Ala623, with a C-terminal 6-His tag Accession # Q9UKQ2
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

A disintegrin and metalloprotease domain-containing protein 28 (ADAM28), also known as MDC-L, is a member of the M12B peptidase family of enzymes. It is synthesized as an approximately 80-90 kDa glycosylated proprotein that is processed to a mature form later in the secretory pathway (1-3). After the removal of the propeptide that contains a cysteine switch motif, the activated form of ADAM28 consists of a 467 aa extracellular domain (ECD) which contains a peptidase, disintegrin, cysteine-rich and EGF-like domains, followed by a 21 aa transmembrane segment and an 89 aa cytoplasmic domain (3). Alternative splicing generates additional isoforms with a variety of substitutions and deletions in the cysteine-rich and EGF-like domains (1). Within the ECD, human ADAM28 shares 73% aa sequence identity with mouse and rat ADAM28 (3). ADAM28 is a cell surface protein that is involved in a variety of cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis (4-6), and exhibits catalytic activity to insulin-like growth factor binding protein-3 (7). Like many members of the ADAM proteins, the disintegrin domain of ADAM28 interacts with integrin to influence cell adhesion and cell-cell interaction (8). ADAM28 can mediate the adhesion of the Jurkat cells, T-lymphoma cell line, through integrin α4β1(9). R&D Systems in-house testing also indicates that ADAM28 cleaves IGFBP-3.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.