

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CEACAM-20 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 1032339
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell HEK293-derived human CEACAM-20 Gln31-Gly450 Accession # Q6UY09-1
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	HEK293 Human Cell Line Transfected with Human CEACAM-20 and eGFP

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Carcinoembryonic antigen-related cell adhesion molecule-20 (CEACAM-20) is a member of the CEACAM subfamily of glycoproteins in the immunoglobulin (Ig) superfamily. Mature human CEACAM-20 consists of a 420 amino acid (aa) extracellular domain (ECD), a 21 aa helical transmembrane segment, and a 114 aa cytoplasmic domain. The extracellular domain possesses four IgC2-like domains which are stabilized by disulfide bonds, as well as several predicted glycosylation sites (1-5). The extracellular domain of CEACAM-20 is also unique among the CEACAMs because it contains a truncated IgV-like N domain (2). Within the ECD, human CEACAM-20 shares 64% and 62% aa identity with the mouse and rat CEACAM-20, respectively. The cytoplasmic domain is unusually long compared to most other CEACAMs and is predicted to contain four tyrosine phosphorylation sites, two of which correspond to the immune-receptor tyrosine-based activation motif (ITAM) (2, 3). Human CEACAM proteins have been linked to numerous intercellular-adhesion and intracellular signaling processes including cell adhesion, growth, and recognition, differentiation, angiogenesis, and apoptosis (7, 8). Human CEACAM-20 expression is limited to the reproductive system and the intestinal tract, with the highest levels of expression found in the small intestine and prostate (2, 3). An *in vitro* model of human prostate morphogenesis showed that CEACAM-20 is co-expressed with CEACAM-1 and plays a critical role in the formation of prostate organoids, making it a marker for prostate cancer (2). Although the exact mechanism is not fully understood, CEACAM-20 may promote the proliferation of intestinal epithelial cells (IECs) (9). There is evidence suggesting CEACAM-20 can induce the production of chemokines like interleukin (IL)-8 and stimulate inflammatory responses in colitis and Crohn's disease (6). CEACAM-20 is also thought to act as a physiological substrate for SAP-1 in the intestinal epithelium (10).

References:

1. Tchoupa, A. *et al.* (2014) J Cell Commun Signal **12**:27.
2. Zhang, H. *et al.* (2013) PLoS ONE **8**:e53359.
3. Zebhauser, R. *et al.* (2005) Genomics **86**:566.
4. Beauchemin, N. Arabzadeh, A. (2013) Cancer Metastasis Rev **32**(3):643.
5. Kuespert, K. *et al.* (2006) Curr Opin Cell Biol **18**:565.
6. Murata, Y. *et al.* (2015) PNAS **E4264**.
7. Obrink, B. (1997) Curr Opin Cell Biol **9**:616.
8. Horst, AK. Wagener, C. (2004) Handb Exp Pharmacol **283**.
9. Kitamura, Y. *et al.* (2015) Genes to Cells **20**:578.
10. Kotani, T. *et al.* (2016) Expert Review of Gastroenterology & Hepatology. **10**:1313.

Human CEACAM-20 Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 1032339

Catalog Number: FAB9416G

100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.