

DESCRIPTION

Species Reactivity	Human
Specificity	Detects Irisin peptide in direct ELISA.
Source	Monoclonal Mouse IgG _{2A} Clone # 1056111
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Synthetic peptide covering the C-term of the Irisin peptide sequences of the FNDC5 gene. Glu97-Glu143 Accession # Q8NAU1
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Irisin (also known as FNDC5) is a 12 kDa glycosylated polypeptide hormone that regulates energy metabolism, stem cell differentiation, and neuronal development (1, 2). Human Irisin is synthesized as a 212 amino acid (aa) precursor encoding a type 1 transmembrane protein with a 121 aa extracellular domain (ECD), a 21 aa transmembrane domain, and a 39 aa cytoplasmic domain. The ECD of Irisin contains a fibronectin type III domain and multiple glycosylation sites. The ECD is proteolytically cleaved to release the 112 aa soluble Irisin hormone into circulation (2-5). Mature human, mouse, and rat Irisin share 100% sequence identity. Expression of Irisin is induced in skeletal muscle and subcutaneous adipose tissue during and shortly after exercise (2, 6). Irisin induces expression of peroxisome proliferator-activated receptor gamma co-activator 1 alpha (PGC1 alpha) and uncoupling protein-1 (UCP1), mitochondrial-associated metabolic proteins (7, 8). Irisin induces the transition of white adipose tissue into more metabolically active beige adipose tissue. In mice, expression of Irisin has been shown to regulate obesity and diabetes (1, 2). A similar function in humans is suggested (9). Irisin also regulates neuronal cell differentiation and neurite outgrowth in the brain and is involved in the differentiation of osteoblasts (10-14).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.