

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Hepcidin in direct ELISA.
Source	Monoclonal Rat IgG _{2A} Clone # 968001
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Synthetic peptide containing mouse Hepcidin Accession # Q9EQ21
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Hepcidin, also known as Liver Expressed Antimicrobial Protein 1 (LEAP-1), is a peptide hormone that is involved in the regulation of iron metabolism (1, 2). It is synthesized as a prohormone that is cleaved intracellularly and secreted as a mature 25 amino acid peptide (1, 3, 4). Hepcidin contains eight cysteine residues that form four disulfide bonds which appear to be important for stability in biological fluids (5). It is predominantly expressed, processed, and secreted by hepatocytes (2, 6). Hepcidin expression is positively regulated by inflammation via IL-6/JAK2/ STAT3 signaling, endoplasmic reticulum stress, and BMP-6 (7-11). BMP-6-dependent Hepcidin induction involves RGM-C/Hemojuvelin, which acts as a co-receptor for BMP-6 (11-13). Conversely, Hepcidin expression is negatively regulated by MMP-15/MT2-MMP and multiple erythropoietic stimuli, including anemia, hypoxia, and Erythropoietin (14-18). MMP-15 downregulates Hepcidin expression by interacting with and cleaving RGM-C (19). Hepcidin was originally identified in human blood and urine as an antimicrobial peptide (1, 3). It has since been shown to regulate iron metabolism. Hepcidin binds the cellular iron exporter Ferroportin, and this interaction results in Ubiquitin-mediated degradation of both Hepcidin and Ferroportin (20-22). Degradation of Ferroportin results in reduced iron release from macrophages, hepatocytes, and duodenal enterocytes, suggesting that Hepcidin may be an effector of inflammatory hypoferrremia (20).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.