

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Acetyl Histone H4 when acetylated at Lys5 in Western blots.
Source	Monoclonal Rabbit IgG Clone # 2129F
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Human Histone H4 synthetic peptide Accession # P62805 (H4_Human)
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Histone4 (H4/HIST1H4I; also H4M and HIST4H4) is a 12-15 kDa member of the histone H4 family of molecules. Human histone 4 is 103 amino acids (aa) in length. Full length histone 4 is identical in aa sequence to mouse histone 4. H4 is ubiquitously expressed. Histones are the organizing core of nucleosomes, forming an octamer comprised of four histone dimers. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries, mediating transcription regulation, DNA repair, DNA replication and chromosomal stability. Acetylation of the Histone H4 lysine residue at position 5 (K5) reduces the positive charge on the histones, relaxing the nucleosome; the relaxed state is associated with higher levels of gene transcription. Misregulation of acetylation has been linked to developmental and neurodegenerative disorders including Alzheimer's disease, and cancer. Butyryl-CoA has been shown to induce the K5 acetylation on H4.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.