

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human 20S Proteasome in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 863425
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human 20S Proteasome purified from human erythrocytes
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The 20S Immunoproteasome is a modified form of the constitutively active 20S Proteasome core particle and is the catalytic subunit of the multi-complex Immunoproteasome. The structure of the 20S Immunoproteasome is similar to the 20S Proteasome, which is composed of 28 non-identical subunits arranged into four stacked rings. However, during 20S Immunoproteasome assembly, the three catalytic beta subunits, beta 1, 2, and 5, in the two interior rings of the 20S Proteasome are replaced by three IFN-gamma-inducible catalytic subunits: beta 1i/LMP2, beta 2i/LMP7, and beta 5i/MECL-1. The 20S Immunoproteasome is commonly associated with the 19S, PA28 alpha/beta, or the PA28 gamma regulatory complexes. 20S Immunoproteasome expression is enriched in antigen presenting cells of the immune system where the 20S Immunoproteasome selectively degrades intracellular proteins in a manner that optimizes the generation of peptides for MHC class I antigen presentation.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.