

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Chorionic Gonadotropin α/β (HCG) and Human CG α in direct ELISAs.
Source	Monoclonal Rabbit IgG Clone # 1177B
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Chinese Hamster Ovary cell line, CHO-derived human Chorionic Gonadotropin α/β (HCG) and Human CG α Ser21-Gln165 and Ala25-Ser116 Accession # P01233 and P01215
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

HCG (human chorionic gonadotropin) is a member of the glycoprotein hormone (GPH) family within the cystine knot growth factor superfamily (1-5). It is a heterodimer of a 23-32 kDa unique subunit, CG β , with a 14-22 kDa alpha subunit, CG α (common glycoprotein hormone alpha) that is shared with GPH family members lutropin (LH), follitropin (FSH) and thyroid stimulating hormone (TSH) (1, 2). CG β occurs only in higher primates, while the most closely related hormone, LH β , is expressed in all mammals (6). Human CG β and LH β share a receptor, LH/CG-R or LHR, and show 86% aa sequence identity between aa 21 and 133, before diverging into a 32 aa, highly O-glycosylated (CG β) or 8 aa (LH β) C-terminal tail (2). Mature human CG α shares 69%-73% aa identity with dog, rabbit, rat, mouse, cow, sheep, pig, cat and horse CG α . Each subunit form a cystine knot structure with three disulfide bridges (5). A "seat-belt" loop of CG β wraps around CG α , stabilizing subunit non-covalent association and conferring receptor selectivity (5). CG β is encoded by six clustered, nonallelic genes that encode identical, but differentially expressed, proteins (2, 7). HCG produced by cytotrophoblast cells in early pregnancy is hyperglycosylated and sialylated, increasing its acidity and half-life (3, 4, 8). Forms with lower glycosylation are produced by syncytiotrophoblasts in continuing pregnancy, and in small amounts by the pituitary where it is also sulfated (4, 8). Free, variably glycosylated CG β subunits are also reported (3, 4). The primary role of HCG is to act as an autocrine factor to establish pregnancy and control placental growth and function. HCG has also been shown to induce the angiogenic factor, EG-VEGF/PK1, and contribute to immune privilege by increasing circulating regulatory T cells and anti-inflammatory cytokines IL-10 and IL-27, via cAMP signaling (9, 10). In addition to pregnancy, large amounts of HCG are produced in gestational trophoblastic diseases such as choriocarcinoma and hydatiform mole (3, 4). HCG may also be produced by ovarian and testicular germ cell tumors and advanced cancers that have dedifferentiated (3, 4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.