

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Angiopoietin-like Protein 4/ANGPTL4 in direct Elisa.
Source	Monoclonal Rat IgG _{2A} Clone # 547113
Purification	Protein A or G purified from cell culture supernatant
Immunogen	CHO-derived mouse Angiopoietin-like Protein 4/ANGPTL4 protein Gln24-Ser410 Accession # Q9Z1P8
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Angiopoietin-like 4 (ANGPTL4), also known as FIAF, FARP, and PGAR, is a 55 kDa glycoprotein secreted by the liver and fat tissue. It is structurally related to the angiopoietins and contains an N-terminal coiled coil domain and a C-terminal fibrinogen-like domain which can be proteolytically separated in vivo (1). Mature mouse ANGPTL4 shares 26%-30% amino acid (aa) sequence identity with mouse ANGPTL1, 2, 3, 4, 6, and 7. It shares 75% and 97% aa sequence identity with human and rat ANGPTL4, respectively. The coiled coil domain, which is not glycosylated, mediates the formation of variable sized disulfide-linked oligomers (2). This domain directly inhibits lipoprotein lipase, resulting in increased circulating triglyceride levels (3, 4). In humans, the N-terminal fragment and full length ANGPTL4 physically associate with HDL (4). In mouse, however, full length ANGPTL4 associates with HDL, while the N-terminal fragment associates with LDL (4). Circulating ANGPTL4 is decreased in type II diabetics with a subsequent loss of its normal plasma glucose lowering activity (5). Its expression in adipose tissue is induced by fasting and suppressed by feeding (6). In hypoxic areas, ANGPTL4 is induced in both vascular endothelial cells and tumor cells (7, 8). The N-terminal fragment can function as an angiogenesis inhibitor (7, 8). In contrast, the C-terminal fragment modulates cell adhesion through interactions with heparan sulfate proteoglycans, Integrins β 1 and β 5, Vitronectin, and Fibronectin, thereby promoting keratinocyte migration and wound healing (7, 9, 10). ANGPTL4 additionally enhances the survival of hematopoietic and mesenchymal stem cells (11, 12). The expression of an undersialylated form of ANGPTL4 in renal podocytes contributes to proteinuria and nephrotic syndrome (13).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.