

DESCRIPTION




Species Reactivity	Goat
Specificity	Detects goat IgG heavy and light chains in direct ELISAs and Western blots. In direct ELISAs, approximately 5% cross-reactivity with human IgG, rabbit IgG, mouse IgG and less than 1% cross-reactivity with chicken IgY is observed.
Source	Polyclonal Donkey IgG
Purification	Antigen Affinity-purified
Immunogen	Goat IgG
Formulation	Supplied as a 0.2 um filtered solution in PBS with stabilizers. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Use at a 1:1000 dilution.
Simple Western	Use at a 1:50 dilution.

DATA

<p>Western Blot</p>  <p>Detection of Goat IgG Primary Antibody by Western Blot Western blot shows detection of Goat Anti-Human/Mouse/Rat GRP75/HSPA9B Polyclonal Antibody (Catalog # AF3584) using a 1:1000 dilution of HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF109). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 2.</p>	<p>Simple Western</p>  <p>Detection of Goat IgG Primary Antibody by Simple Western™. Simple Western lane view shows detection of Goat Anti-Human/Mouse/Rat TC-PTP Polyclonal Antibody (Catalog # AF1930) using a 1:50 dilution of HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF109). This experiment was conducted under reducing conditions and using the 12-230 kDa separation system.</p> 
---	--

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Do not freeze. <ul style="list-style-type: none"> 6 months from date of receipt, 2 to 8 °C as supplied.