

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-4 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2A} Clone # 1067629
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese Hamster Ovary cell line, CHO-derived human IL-4 His25-Ser153 Accession # P05112
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Intracellular Staining by Flow Cytometry	Titration recommended for optimal concentration with starting range of 0.1-1 µg/1 million cells. Sample used for this experiment was PBMCs lymphocytes.
---	---

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Use a manual defrost freezer and avoid repeated freeze-thaw cycles. • 12 months, 2 to 8 °C under sterile conditions after opening.

BACKGROUND

Interleukin-4 (IL-4), also known as B cell-stimulatory factor-1, is a monomeric, approximately 13 kDa-18 kDa Th2 cytokine that shows pleiotropic effects during immune responses (1-3). It is a glycosylated polypeptide that contains three intrachain disulfide bridges and adopts a bundled four α-helix structure (4). Human IL-4 is synthesized with a 24 aa signal sequence. Alternate splicing generates an isoform with a 16 aa internal deletion. Mature human IL-4 shares 55%, 39% and 43% aa sequence identity with bovine, mouse, and rat IL-4, respectively. Human, mouse, and rat IL-4 are species-specific in their activities (5-7). IL-4 exerts its effects through two receptor complexes (8, 9). The type I receptor, which is expressed on hematopoietic cells, is a heterodimer of the ligand binding IL-4 Rα and the common γ chain (a shared subunit of the receptors for IL-2, -7, -9, -15, and -21). The type II receptor on nonhematopoietic cells consists of IL-4 Rα and IL-13 Rα1. The type II receptor also transduces IL-13 mediated signals. IL-4 is primarily expressed by Th2-biased CD4+ T cells, mast cells, basophils, and eosinophils (1, 2). It promotes cell proliferation, survival, and immunoglobulin class switch to IgG4 and IgE in human B cells, acquisition of the Th2 phenotype by naïve CD4+ T cells, priming and chemotaxis of mast cells, eosinophils, and basophils, and the proliferation and activation of epithelial cells (10-13). IL-4 plays a dominant role in the development of allergic inflammation and asthma (12, 14).

References:

1. Benczik, M. and S.L. Gaffen (2004) Immunol. Invest. **33**:109.
2. Chomarat, P. and J. Banchereau (1998) Int. Rev. Immunol. **17**:1.
3. Yokota, T. *et al.* (1986) Proc. Natl. Acad. Sci. **83**:5894.
4. Redfield, C. *et al.* (1991) Biochemistry **30**:11029.
5. Ramirez, F. *et al.* (1988) J. Immunol. Meth. **221**:141.
6. Leitenberg, D. and T.L. Feldbush (1988) Cell. Immunol. **111**:451.
7. Mosman, T.R. *et al.* (1987) J. Immunol. **138**:1813.
8. Mueller, T.D. *et al.* (2002) Biochim. Biophys. Acta **1592**:237.
9. Nelms, K. *et al.* (1999) Annu. Rev. Immunol. **17**:701.
10. Paludan, S.R. (1998) Scand. J. Immunol. **48**:459.
11. Corthay, A. (2006) Scand. J. Immunol. **64**:93.
12. Ryan, J.J. *et al.* (2007) Crit. Rev. Immunol. **27**:15.
13. Grone, A. (2002) Vet. Immunol. Immunopathol. **88**:1.
14. Rosenberg, H.F. *et al.* (2007) J. Allergy Clin. Immunol. **119**:1303.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.