

## Human SF20/MYDGF Alexa Fluor® 350-conjugated Antibody

Monoclonal Mouse IgG<sub>1</sub> Clone # 1009420

Catalog Number: IC1147U

100 µg

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human SF20 in direct ELISAs.		
Source	Monoclonal Mouse IgG <sub>1</sub> Clone # 1009420		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human SF20 Met1-Leu173 Accession # Q969H8		
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm		
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS				
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.				
	Recommended Concentration	Sample		
Intracellular Staining by Flow Cytometry	0.25-1 μg/10 <sup>6</sup> cells	U937 Human Histiocytic Lymphoma Cell Line fixed with paraformaldehyde and permeabilized with saponin		

PREPARATION AND STORAGE			
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.		
Stability & Storage	Protect from light. Do not freeze.		
	12 months from date of receipt, 2 to 8 °C as supplied.		

## BACKGROUND

Human SF20 (Stromal cell-derived growth factor SF20, C19orf10, MYDGF1, Myeloid-derived growth factor1 also formerly known as IL25) is a 15 kDa (predicted) bone marrow-derived monocyte and paracrine-acting protein that promotes cardiac myocyte survival and adaptive angiogenesis for cardiac protection and/or repair after myocardial infarction. SF20 stimulates endothelial cell proliferation through a MAPK1/3-, STAT3- and CCND1 mediated signaling pathway. It is thought to inhibit cardiac myocyte apoptosis in a PI3K/AKT-dependent signaling pathway and is involved in endothelial cell proliferation and angiogenesis. Human SF20 shares 87% aa identity with mouse SF20.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

