

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human TLR3 in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) TLR1, rhTLR2, rhTLR4 and recombinant mouse TLR6 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human TLR3 Lys27-Ser711 Accession # O15455
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Human TLR3 is a 116 kDa type I transmembrane glycoprotein that belongs to the mammalian Toll-Like Receptor family of pathogen pattern recognition molecules (1, 2). There are at least eleven mouse and ten human members that activate the innate immune system following exposure to a variety of microbial species (3). The human TLR3 cDNA encodes a 904 amino acid (aa) precursor that contains a 23 aa signal sequence, a 681 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 179 aa cytoplasmic region (4). The horseshoe shaped ECD (5, 6) contains 23 leucine rich repeats, and the cytoplasmic domain contains one Toll/IL-1 Receptor (TIR) domain. The ECD of human TLR3 shares 80%, 79%, and 77% aa sequence identity with the ECD of rat, mouse, and bovine TLR3, respectively. TLR3 is found in phagosomes (7), where the acidic pH enables binding of internalized double stranded RNA and mRNA from viruses, parasites, and necrotic virally-infected cells (8-11). Ligand binding by TLR3 induces receptor dimerization (5, 6, 8) leading to the release of inflammatory cytokines and dendritic cell maturation (9, 11-13). TLR3 is expressed in dendritic cells, macrophages, microglia, and astrocytes (13-15) and is upregulated by IFN- β and LPS (9, 14). TLR3 expression is also induced by lung fibroblasts and epithelial cells by respiratory syncytial virus infection (12).

References:

1. Sen, G.C. and S.N. Sarkar (2005) Cytokine Growth Factor Rev. **16**:1.
2. Schroder, M. and A.G. Bowie (2005) Trends Immunol. **26**:462.
3. Hopkins, P.A. and S. Sriskandan (2005) Clin. Exp. Immunol. **140**:395.
4. Rock, F.L. *et al.* (1998) Proc. Natl. Acad. Sci. **95**:588.
5. Choe, J. *et al.* (2005) Science **309**:581.
6. Bell, J.K. *et al.* (2005) Proc. Natl. Acad. Sci. **102**:10976.
7. Nishiya, T. *et al.* (2002) J. Biol. Chem. **280**:37107.
8. de Bouteiller, O. *et al.* (2005) J. Biol. Chem. **280**:38133.
9. Alexopoulou, L. *et al.* (2001) Nature **413**:732.
10. Aksoy, E. *et al.* (2005) J. Biol. Chem. **280**:277.
11. Kariko, K. *et al.* (2004) J. Biol. Chem. **279**:12542.
12. Rudd, B.D. *et al.* (2005) J. Virol. **79**:3350.
13. Farina, C. *et al.* (2005) J. Neuroimmunol. **159**:12.
14. Heinz, S. *et al.* (2003) J. Biol. Chem. **278**:21502.
15. Wang, T. *et al.* (2004) Nat. Med. **10**:1366.