

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human and mouse SOX2 in Western blots.
Source	Monoclonal Mouse IgG _{2A} Clone # 245610
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human SOX2 Gly135-Met317 Accession # P48431
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

SOX2 belongs to the SOX (SRY-like HMG box) family of transcription factors with diverse roles in development. SOX2 functions in specifying the first three lineages present at implantation and in regulating proliferation and differentiation in the developing peripheral nervous system (1-5).

References:

- Graham, V. *et al.* (2003) *Neuron* **39**:749.
- Avilion, A.A. *et al.* (2003) *Genes Dev.* **17**:126.
- Kishi, M. *et al.* (2000) *Development* **127**:791.
- Yuan, H. *et al.* (1995) *Genes Dev.* **9**:2635.
- Uwanogho, D. *et al.* (1995) *Mech. Dev.* **49**:23.
- Stevanovic, M. (2003) *Mol. Biol. Rep.* **30**:127.