

Human Langerin/CD207 Alexa Fluor® 647-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 343828

Catalog Number: IC2088R

100 µg

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Langerin/CD207 in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 343828
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Langerin/CD207 Tyr64-Pro328 Accession # Q9UJ71
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human immature dendritic cells differentiated from human peripheral blood mononuclear cell derived CD14 ⁺ cells treated with 20 ng/mL Recombinant Human IL-4 (Catalog # 204-IL) and 50 ng/mL Recombinant Human GM-CSF (Catalog # 215-GM) for 6-7 days

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Langerin (also known as CD207) is a type II transmembrane glycoprotein which is member K of the C-type lectin domain family 4 (1). Langerin is used as a marker for Langerhans cells (LCs) which represent the immature dendritic cells in the epidermis (1, 2). LCs uniquely contain "tennis racket"-shaped endosomal recycling compartment subdomains with pentalamellar membranes termed Birbeck granules (1-3). Langerin is necessary and sufficient for Birbeck granule formation (1). The 328 amino acid (aa) human langerin sequence contains a 43 aa cytoplasmic domain, a 21 aa transmembrane domain and a 264 aa extracellular domain (ECD) that contains a coiled-coil domain and a single C-type lectin domain. Trimerization greatly increases the lectin binding affinity (4). Langerin internalizes endogenous proteins such as type I procollagen. Internalization by LC is thought to lead to suppression of self reactions (4-6). Langerin also mediates endocytosis of non-peptide antigens containing mannose, N-acetyl glucosamine and fucose that are expressed by mycobacteria and fungi (4, 7). Some antigens, such as the *M. leprae* glycolipid arabinomycolate, are ultimately presented by human LC CD1a in cutaneous-draining lymph nodes (8). Langerin performs a barrier-like function to HIV-1 transmission due to its internalization of virus particles for destruction (9). A rare human polymorphism within the lectin domain, W264R, abolishes both carbohydrate recognition and Birbeck granule formation (10, 11). Genetic deletion of mouse langerin was not shown to have functional consequence other than abolishing Birbeck granule formation (12). Human langerin shares 68%, 62%, 71% aa identity with mouse, rat and bovine langerin ECD, respectively.

References:

- Valladeau, J. *et al.* (2000) *Immunity* **12**:71.
- Valladeau, J. *et al.* (2003) *Immunol. Res.* **28**:93.
- McDermott, R. *et al.* (2002) *Mol. Biol. Cell* **13**:317.
- Stambach, N. S. and M. E. Taylor (2003) *Glycobiology* **13**:401.
- Tada, Y. *et al.* (2006) *J. Invest. Dermatol.* **126**:1549.
- Ritter, U. and A. Osterloh (2007) *Med. Microbiol. Immunol.* **196**:51.
- Takahara, K. *et al.* (2003) *Int. Immunol.* **16**:819.
- Hunger, R. E. *et al.* (2004) *J. Clin. Invest.* **113**:701.
- De Witte, L. *et al.* (2007) *Nat. Med.* **13**:367.
- Verdijk, P. *et al.* (2005) *J. Invest. Dermatol.* **124**:714.
- Ward, E. M. *et al.* (2006) *J. Biol. Chem.* **281**:15450.
- Kissenpfennig, A. *et al.* (2005) *Mol. Cell. Biol.* **25**:88.

Human Langerin/CD207 Alexa Fluor® 647-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 343828

Catalog Number: IC2088R
100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.