

DESCRIPTION


Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat GSK-3 β in ELISAs and Western blots. Does not cross-react with recombinant human GSK-3 α .
Source	Monoclonal Rat IgG _{2A} Clone # 272536
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human GSK-3 β Met1-Thr420 Accession # P49841
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Glycogen Synthase Kinase-3 (GSK-3) is a serine/threonine kinase initially identified as an inhibitor of glycogen synthase. Two isoforms (GSK-3 α and GSK-3 β) share 85% amino acid sequence identity. GSK-3 β , inhibited by phosphorylation at S9 by Akt, is involved in energy metabolism, body pattern formation, and neuronal cell development. Depending upon the substrate, phosphorylation may either promote activity (SIK-1; OGT; ZC3HAV1) or downregulate activity (SNAI1; BMAL1; MARK2).