

DESCRIPTION



Species Reactivity	Mouse
Specificity	Detects mouse Nanog in Western blots. In this format, approximately 50% cross-reactivity with recombinant human Nanog is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Nanog Trp154-Leu262 Accession # Q80Z64
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunocytochemistry	8-25 µg/mL	See Below
Intracellular Staining by Flow Cytometry	5 µL/10 ⁶ cells	See Below

DATA

<p>Intracellular Staining by Flow Cytometry</p>  <p>Detection of Nanog in D3 Mouse Cell Line by Flow Cytometry. D3 mouse embryonic stem cell line either untreated (dark orange filled histogram) or treated with retinoic acid for 3 days (light orange filled histogram) was stained with Goat Anti-Mouse Nanog Alexa Fluor® 488-conjugated Antigen Affinity-purified Polyclonal Antibody (Catalog # IC2729G) or isotype control antibody (Catalog # IC108G, blue open histogram). To facilitate intracellular staining, cells were fixed with Flow Cytometry Fixation Buffer (Catalog # FC004) and permeabilized with Flow Cytometry Permeabilization/Wash Buffer I (Catalog # FC005). View our protocol for Staining Intracellular Molecules.</p>	<p>Immunocytochemistry</p>  <p>Nanog in D3 Mouse Embryonic Stem Cells. Nanog was detected in immersion fixed D3 mouse embryonic stem cell line on irradiated mouse embryonic fibroblasts using Goat Anti-Mouse Nanog Alexa Fluor® 488-conjugated Antigen Affinity-purified Polyclonal Antibody (green; Catalog # IC2729G) at 10 µg/mL for 3 hours at room temperature. Cells were counterstained with DAPI (blue). Specific staining was localized to nuclei. View our protocol for Fluorescent ICC Staining of Stem Cells on Coverslips.</p>
---	--

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Nanog is a member of the homeobox family of DNA binding transcription factors that has been shown to maintain pluripotency of embryonic stem cells. Its expression is high in undifferentiated embryonic stem cells and is down-regulated during embryonic stem cell differentiation, concomitant with loss of pluripotency (1-3).

References:

1. Mitsui, K. *et al.* (2003) *Cell* 11:631.
2. Chambers, I. *et al.* (2003) *Cell* 113:643.
3. Hart, A.H. *et al.* (2004) *Dev. Dyn.* 230:187.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc. and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.