

DESCRIPTION

| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects human STAT1 when phosphorylated at Y701 in Western blots. |
| Source | Recombinant Monoclonal Rabbit IgG Clone # 1086B |
| Purification | Protein A or G purified from cell culture supernatant |
| Immunogen | Phosphopeptide containing the human STAT1 Y701 site Accession # NP009330 |
| Conjugate | Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|---|----------------------------------|---|
| Intracellular Staining by Flow Cytometry | 0.25-1 µg/10 ⁶ cells | IFN alpha-treated Daudi Human Cell Line fixed with paraformaldehyde and permeabilized with methanol |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

STAT1 (signal transducer and activator of transcription #1) is an 88 kDa member of the STAT family of cytoplasmic transcription factors. STAT members generally mediate cytokine, growth factor and hormone receptor signal transduction. STAT1 is associated with type I and II interferon signaling. All STATs contain an N-terminal oligomerization domain, a DNA-binding domain, and an SH2-association region. STAT1 is phosphorylated at Y701 by receptor-associated Janus kinases (JAKs) leading to STAT1 dimerization and subsequent translocation to the nucleus to activate gene transcription.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.