

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CCL13/MCP-4 in ELISAs. In sandwich immunoassays, less than 1% cross-reactivity with recombinant human (rh) CCL2 is observed and no significant cross-reactivity or interference with rhCCL5, 7, 8, 11, 24, recombinant mouse CCL2, 5, 7, 11 or 12 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 73506
Purification	Protein A or G purified from ascites
Immunogen	<i>E. coli</i> -derived recombinant human CCL13/MCP-4 Gln24-Thr98 Accession # Q99616.1
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	A549 human lung carcinoma cell line treated with recombinant human IL-1β, fixed with paraformaldehyde and permeabilized with saponin

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CCL13, also known as Monocyte Chemoattractant Protein-4 (MCP-4), is a CC chemokine that acts as a chemoattractant for monocytes, eosinophils and T cells and as an activator of basophils. Human CCL13 cDNA encodes a 98 amino acid residue precursor protein with a 23 amino acid residue hydrophobic signal peptide that is cleaved to yield an 8 kDa, 75 aa mature CCL13. Mature CCL13 lacks any potential N-glycosylation sites and shares a pyroglutamate proline motif with other human MCP proteins. Human CCL13 is most homologous to MCP-1, 3 and Eotaxin, exhibiting approximately 65-66% amino acid sequence identity. CCL13 mRNA is expressed by a number of activated cell types, including endothelial cells, macrophages, bronchial epithelium and type II alveolar cells, and perhaps lymphocytes. The bioactivities of CCL13 are mediated by the CC chemokine receptors CCR-2 and CCR-3.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.