

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human Calreticulin in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 326203
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Calreticulin Met1-Asn180 Accession # P27797
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Human Calreticulin is a 55-60 kDa, 400 amino acid, variably glycosylated intra- and extracellular Ca⁺⁺-binding lectin that is ubiquitously expressed. It consists of three domains: a 180 aa N-terminal globular region, a 111 aa P-, or proline rich domain, and a 109 aa C-terminus. The 180 aa N-terminus (aa 18-197) is termed Vasostatin. It is unclear if it is ever generated naturally via proteolytic processing. Vasostatin domain has many functions. It binds to RNA (aa 18-27), has autocatalytic phosphorylase activity (aa 77-197), binds to a KxFFKR motif on steroid hormone receptors, and serves as a lectin-type chaperone for ER-localized molecules. It also shows antiangiogenic activity, presumably by binding to laminin carbohydrates and blocking endothelial cell adhesion and proliferation. Human Calreticulin is 94% aa identical to mouse and rat Calreticulin.