

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human MYF-5 in direct ELISAs.
Source	Monoclonal Mouse IgG _{2A} Clone # 593128
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human MYF-5 Met1-Leu255 Accession # P13349
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	C2C12 mouse myoblast cell line fixed with paraformaldehyde and permeabilized with saponin

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

MYF-5 (Myogenic Factor 5), also known as bHLHc2, is a 37-39 kDa transcriptional activator that belongs to the bHLH superfamily of myogenic regulatory factors. It is found in embryonic skeletal muscle progenitors and neurons, and is induced by SHH and Wnt1. Expression of MYF-5 protein prompts nonmuscle cells to adopt a muscle precursor phenotype. Notably, neurons appear to block MYF-5 accumulation, thus preserving their unique phenotype. MYF-5 forms heterodimers with multiple molecules. When complexed to I-mfa, MYF-5 is retained in the cytoplasm and is inactive; when complexed to E12, MYF-5 is transcriptionally active. Phosphorylation regulates MYF-5 stability. Human MYF-5 is 255 amino acids (aa) in length. It contains one myogenic basic region (aa 1-83) followed by an HLH motif (aa 96-135). Full-length human MYF-5 (aa 1-255) shares 89% aa sequence identity with mouse MYF-5.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.