

Human STAT5a/b Alexa Fluor® 647-conjugated Antibody

Recombinant Monoclonal Rabbit IgG Clone # 1247C Catalog Number: IC41901R

100 µg

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human STAT5a/b when phosphorylated at Y964/Y699 in direct ELISAs and Western blots.		
Source	Recombinant Monoclonal Rabbit IgG Clone # 1247C		
Purification	Protein A or G purified from cell culture supernatant		
Immunogen	Phosphopeptide containing human STAT5b Y699 site (amino acid sequence of this peptide is identical to a corresponding region of human STAT5a containing Y694) Accession # P51692		
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm		
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS				
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.				
	Recommended Concentration	Sample		
Intracellular Staining by Flow Cytometry	0.25-1 μg/10 ⁶ cells	IFN alpha-treated Daudi Human Cell Line fixed with paraformaldehyde and permeabilized with methanol		

PREPARATION AND STORAGE			
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.		
Stability & Storage	Protect from light. Do not freeze.		
	 12 months from date of receipt, 2 to 8 °C as supplied. 		

BACKGROUND

Signal transduction and activator of transcription 5 (STAT5) is a member of the Jak/STAT signal transduction pathway and is activated by a variety of cytokines (IL-22, IL-6, IFN-α). STAT5 has two isoforms (A and B) that share 93% amino acid identity and bind the DNA consensus site TTCN₃GAA. STAT5 mediates cytokine signaling by acting as a signal transducer in the cytoplasm and, upon phosphorylation, translocates to the nucleus and activates transcription of specific genes. STAT5 is involved in a wide array of biological processes ranging from regulating apoptosis to adult mammary gland proliferation, differentiation, and survival.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

