


DESCRIPTION	
<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human IRF5 in Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human IRF5 Ala214-Thr395 Accession # NP_002191
<b>Conjugate</b>	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

IRF5 is a member of the IRF family of transcription factors, a family characterized by a helix-turn-helix DNA binding domain enriched in tryptophan repeats. IRF family members show diverse cellular regulation of interferon-stimulated gene transcription, viral-mediated gene activation, apoptosis, differentiation, and cellular growth. IRF5, along with IRF7, are the key mediators of TLR signaling. IRF5 forms heterodimers with IRF3 both of which are necessary for interferon gene transcription. IRF5 knock out mice indicate that IRF5 is critical for induction of apoptosis.