

Human/Mouse RelA/NFκB p65 Alexa Fluor® 750-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 532301

Catalog Number: IC5078S

100 µg

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse RelA/NFκB p65 in Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 532301
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human RelA/NFκB p65 isoform 1 Asn456-Ser551 Accession # Q04206
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	HeLa human cervical epithelial carcinoma cell line, fixed with paraformaldehyde and permeabilized with methanol

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

RelA p65 (v-rel reticuloendotheliosis viral oncogene homolog A) is a 65 kDa member of the NFκB family of nuclear transcription factors. Dimers of p65 with the p50 subunit are the most common form of the NFκB transcription factor, but dimers with it or other family members can also occur. Upon activation, RelA p65 forms a heterotetramer and moves into the nucleus where it binds to specific DNA sequences. An alternatively spliced isoform that lacks amino acids (aa) 222-231 (p65Δ) does not bind DNA. Over the sequence used as an immunogen, human RelA p65 shares 96% and 98% aa identity with mouse and rat RelA p65, respectively. This portion includes one of eight potential ser/thr phosphorylation sites, two acetylation sites, and most of the Rel homology domain that interacts with IκB inhibitors.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.