

## Human Indoleamine 2,3-dioxygenase/IDO Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG2B Clone # 998736 Catalog Number: IC60302G

100 µg

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human Indoleamine 2,3-dioxygenase/IDO in direct ELISAs.		
Source	Monoclonal Mouse IgG <sub>2B</sub> Clone # 998736		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	E. coli-derived human Indoleamine 2,3-dioxygenase/IDO Ala2-Gly403 Accession # P14902		
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm		
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS				
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.				
	Recommended Concentration	Sample		
Flow Cytometry	0.25-1.ug/10 <sup>6</sup> cells	Human PBMC monocytes fixed with 1% paraformaldehyde and permeabilized with saponin		

 $0.25-1 \mu g/10^6 cells$ 

PREPARATION AND STORAGE			
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.		
Stability & Storage	ge Protect from light. Do not freeze.		
	<ul> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>		

Indoleamine 2,3-dioxygenase (IDO) is a heme-containing intracellular dioxygenase catalyzing the degradation of the essential amino acid L-tryptophan to N-formyl-kynurenine (1). This degradation is the first and rate-limiting step of the L-kynurenine pathway (2). IDO is widely expressed in dendritic cells, macrophages, microglia, eosinophils, fibroblasts, endothelial cells, and most tumor cells. In immune cells, its expression is mainly induced by cytokines such as IFN-γ, IFN-α, IFN-β, and IL-10. IDO has an antimicrobial function due to its decreasing the availability of the essential amino acid tryptophan in inflammatory environments (3). Recent studies have demonstrated that IDO induces immunosuppression during infection, pregnancy, transplantation, autoimmunity, and neoplasia

## References:

- 1. Lewis-Ballester, A. et al. (2009) Proc. Natl. Acad. Sci. USA. 106:17371.
- 2. Costantino, G. (2009) Expert Opin. Ther. Targets 13:247.
- 3. Xu, H. et al. (2008) Immunol. Lett. 121:1.
- 4. Lob, S. et al. (2009) Nat. Rev. Cancer 9:445
- 5. Curti, A. et al. (2009) Blood 113:2394.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

biotechne<sup>®</sup>