

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human BPI in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 971526
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell line HEK-293-derived transfected with human BPI Val32-Lys487 Accession # P17213
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	HL-60 Human acute promyelocytic leukemia cell line fixed with Flow Cytometry Fixation Buffer (Catalog # FC004) and permeabilized with Flow Cytometry Permeabilization/Wash Buffer I (Catalog # FC005)

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. • 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Bactericidal/Permeability Increasing protein (BPI) is a 55 kDa antibacterial glycoprotein that plays a role in innate immunity (1, 2). It belongs to the lipid transfer protein family that also includes LPS binding protein (LBP), cholesteryl ester transfer protein (CETP), and phospholipid transfer protein (PLTP). Circulating levels of BPI are positively correlated with the levels of cholesterol, LDL cholesterol, and HDL cholesterol (3). Mature human BPI shares approximately 55% amino acid (aa) sequence identity with mouse and rat BPI. It can be secreted as a monomer or as a disulfide-linked homodimer (4). It consists of a highly basic N-terminal and a hydrophobic C-terminal domain (5). Its N-terminal domain confers the ability of BPI to bind bacterial lipopolysaccharide (LPS) found in the cell walls of Gram negative bacteria and to induce the lysis and phagocytosis of these bacteria (6-9). It also blocks the endothelial cell response to endotoxin (10). BPI is stored in neutrophil and eosinophil granules for induced secretion during inflammation (11, 12). It is additionally expressed in mucosal epithelia and testis (10, 13). BPI can be retained on the surface of both neutrophils and epithelial cells, presumably by its hydrophobic C-terminal domain (8, 10). BPI also functions as an anti-angiogenic molecule by inhibiting vascular endothelial cell proliferation and tubule formation (14). Like the antibacterial actions, this function is mediated by the N-terminal region (15).

References:

- Schultz, H. and J.P. Weiss (2007) Clin. Chim. Acta **384**:12.
- Holweg, A. *et al.* (2011) Biochem. Soc. Trans. **39**:1045.
- Esteve, E. *et al.* (2010) Thromb. Haemost. **103**:780.
- Horwitz, A.H. *et al.* (1996) Protein Exp. Purif. **8**:28.
- Gray, P.W. *et al.* (1989) J. Biol. Chem. **264**:9505.
- Ooi, C.E. *et al.* (1987) J. Biol. Chem. **262**:14891.
- Tobias, P.S. *et al.* (1997) J. Biol. Chem. **272**:18682.
- Weersink, A.J. *et al.* (1993) J. Immunol. **150**:253.
- Nishimura, H. *et al.* (2001) Immunology **103**:519.
- Canny, G. *et al.* (2002) Proc. Natl. Acad. Sci. **99**:3902.
- Weiss, J. and I. Olsson (1987) Blood **69**:652.
- Calafat, J. *et al.* (1998) Blood **91**:4770.
- Lennartsson, A. *et al.* (2005) J. Leukoc. Biol. **77**:369.
- van der Schaft, D.W.J. *et al.* (2000) Blood **96**:176.
- Rauniar, R.K. *et al.* (2002) Invest. Ophthalmol. Vis. Sci. **43**:503.

Human BPI Alexa Fluor® 405-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 971526

Catalog Number: IC74681V

100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.