

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-10 in direct ELISAs.
Source	Recombinant Monoclonal Rabbit IgG _{2B} Clone # 2050B
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human IL-10 Met1-Asn178 Accession # P22301
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human peripheral blood mononuclear cells (PBMCs) stimulated to induce Th2 cells were fixed with Flow Cytometry Fixation Buffer (Catalog # FC004) and permeabilized with methanol

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Interleukin 10, also known as cytokine synthesis inhibitory factor (CSIF), is the charter member of the IL-10 family of α-helical cytokines that also includes IL-19, IL-20, IL-22, IL-24, and IL-26/AK155 (1, 2). IL-10 is secreted by many activated hematopoietic cell types as well as hepatic stellate cells, keratinocytes, and placental cytotrophoblasts (2-5). Mature human IL-10 shares 72%-86% amino acid sequence identity with bovine, canine, equine, feline, mouse, ovine, porcine, and rat IL-10. Whereas human IL-10 is active on mouse cells, mouse IL-10 does not act on human cells (6, 7). IL-10 is a 178 amino acid molecule that contains two intrachain disulfide bridges and is expressed as a 36 kDa noncovalently associated homodimer (6, 8, 9). The IL-10 dimer binds to two IL-10 Rα/IL-10 R1 chains, resulting in recruitment of two IL-10 Rβ/IL-10 R2 chains and activation of a signaling cascade involving JAK1, TYK2, and STAT3 (10). IL-10 Rβ does not bind IL-10 by itself but is required for signal transduction (1). IL-10 Rβ also associates with IL-20 Rα, IL-22 Rα, or IL-28 Rα to form the receptor complexes for IL-22, IL-26, IL-28, and IL-29 (11-13). IL-10 is a critical molecule in the control of viral infections and allergic and autoimmune inflammation (14-16). It promotes phagocytic uptake and Th2 responses but suppresses antigen presentation and Th1 proinflammatory responses (2).

References:

1. Pestka, S. *et al.* (2004) *Annu. Rev. Immunol.* **22**:929.
2. Sabat, R. *et al.* (2010) *Cytokine Growth Factor Rev.* **21**:331.
3. Mathurin, P. *et al.* (2002) *Am. J. Physiol. Gastrointest. Liver Physiol.* **282**:G981.
4. Grewe, M. *et al.* (1995) *J. Invest. Dermatol.* **104**:3.
5. Szonyi, B.J. *et al.* (1999) *Mol. Hum. Reprod.* **5**:1059.
6. Vieira, P. *et al.* (1991) *Proc. Natl. Acad. Sci.* **88**:1172.
7. Hsu, D.-H. *et al.* (1990) *Science* **250**:830.
8. Windsor, W.T. *et al.* (1993) *Biochemistry* **32**:8807.
9. Syto, R. *et al.* (1998) *Biochemistry* **37**:16943.
10. Kotenko, S.V. *et al.* (1997) *EMBO J.* **16**:5894.
11. Kotenko, S.V. *et al.* (2000) *J. Biol. Chem.* **276**:2725.
12. Hor, S. *et al.* (2004) *J. Biol. Chem.* **279**:33343.
13. Sheppard, P. *et al.* (2003) *Nat. Immunol.* **4**:63.
14. Fitzgerald, D.C. *et al.* (2007) *Nat. Immunol.* **8**:1372.
15. Wu, K. *et al.* (2007) *Cell. Mol. Immunol.* **4**:269.
16. Blackburn, S.D. and E.J. Wherry (2007) *Trends Microbiol.* **15**:143.

Human IL-10 Alexa Fluor® 700-conjugated Antibody

Recombinant Monoclonal Rabbit IgG_{2B} Clone # 2050B

Catalog Number: IC9210N

100 µg

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.