
MATERIAL DATA SHEET

Ubi-Detect, Protein Ubiquitination Detection Kit
Cat. # K-410

Covalent attachment of ubiquitin to target proteins is one of the most versatile and interesting post-translational modifications observed in eukaryotic organisms. Because polyubiquitin chains may be generated by connecting adjacent ubiquitin molecules in one of eight different orientations, a large number of varied architectures may be generated. Polyubiquitination of proteins directs them to different cellular fates such as translocation to different areas within the cell, assembly into multi-subunit complexes, or degradation via the proteasome, depending on the type of linkage used in the polyubiquitin chain. It should not be assumed however that polyubiquitination of a protein is required to generate a ubiquitin-dependent biological signaling event. Indeed, monoubiquitination (including multi-monoubiquitination) of many proteins is necessary and sufficient to provide cues that are important in processes such as viral budding, DNA repair, receptor endocytosis, and gene expression. It is increasingly clear that identifying and potentially altering the ubiquitination status of numerous proteins will be important in understanding and treating multiple human disease states.

The Ubi-Detect Protein Ubiquitination Detection Kit is a practical tool for helping the investigator answer a straightforward question: is a protein of interest ubiquitinated? This kit provides reagents and a protocol that may be used to isolate a protein (or protein complex) by immunoprecipitation, and then analyze the complex for the presence of ubiquitin. Samples from cell lysates or other complex mixtures may be analyzed.

NOTE: Kit contains reagents sufficient for preparing 12 lysates from 6-well plates or T25 tissue culture flasks, and 2 Western Blots (small or medium sized gel format).

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565
www.bostonbiochem.com

Technical Hints

It is essential that the investigator read the kit instructions before initiating any experiments.

Ubi-Detect Protein Ubiquitination Detection Kit data output consists of SDS-PAGE gel-based detection of ubiquitin via Western Blot. The ubiquitination status of a protein is potentially influenced by a large number of factors depending on the identity of the protein. Some proteins are ubiquitinated in response to changes in the cell-cycle, activation of cell-surface receptors, DNA damage, etc. It is not in the scope of this protocol to guide the investigator on how and when a protein of interest may be ubiquitinated. It is assumed that a set of experimental conditions has been defined by the investigator in which the (potential) ubiquitination of a protein will be examined.

The reagents and protocol provided in this kit may result in the isolation of protein complexes via immunoprecipitation ("co-IP"). In some cases the protein of interest might not be ubiquitinated, but is part of a co-IP'd multi-protein complex containing one or more ubiquitinated proteins. This could lead to an errant conclusion about the ubiquitination of the target protein. There are multiple techniques to address this issue—contact techsupport@bostonbiochem.com for more details.

Complex lysates from eukaryotic cells contain deubiquitinases ("DUBs") that can quickly remove ubiquitin from a target protein of interest. The DUB-Stop Protease Inhibitor Mix included with this kit contains fast-acting compounds that provide excellent protection against most DUBs. For additional protection, consider adding 2-10 μ M ubiquitin-aldehyde, ubiquitin-vinyl sulfone, ubiquitin-vinyl methyl ester, or ubiquitin-propargyl amide to help prevent unwanted DUB activity.

The ability to detect ubiquitination of low-abundance proteins is often enhanced by inhibiting proteasome activity for 4-12 hours prior to generating cellular lysates. The use of cell-permeable proteasome inhibitors such as epoxomicin may be employed in these instances.

SDS-PAGE gels, PVDF membrane, blocking reagent, and protocols provided or referenced in this kit have been chosen following careful testing at Boston Biochem. Modifications to the protocol or selection of alternative reagents may require assay optimization by the end-user. More information is available at techsupport@bostonbiochem.com.

Materials Provided and Storage Conditions

<u>Component</u>	<u>Volume</u>
1. 10X Ubl Lyse Reagent	2.4 ml
2. 100X DUB-Stop Protease Inhibitor Mix	240 µl
3. 5X HS Wash Buffer	4.8 ml
4. 10X LS Wash Buffer	1.2 ml
5. 100X α-ubiquitin antibody, biotinylated	80 µl
6. Ubiquitinated Positive Control Sample	36 µl

Storage: Store components at -20°C. Avoid multiple freeze/thaw cycles.

Other Materials Required

The investigator will need to obtain the following reagents prior to using this kit.

dH ₂ O	Sterile
SDS-PAGE Gels	Refer to Protocols for guidance.
SDS-PAGE Sample Buffer	Compatible with SDS-PAGE gel.
Dithiothreitol (DTT)	1M in dH ₂ O (Pierce #20290) or similar
IP antibody	User-validated antibody used to immunoprecipitate ("IP") target protein of interest
PBST	1X PBS (Calbiochem #524650) + 0.05% Tween-20 (Sigma #P1379) or similar
BSA	OmniPur BSA (EMD-Millipore #2910) or similar
Towbin Buffer	25 mM Tris Base, 192 mM glycine, 20% methanol (pH ~ 8.3. Do not adjust pH)
PVDF Membrane	Immun-Blot PVDF Membrane (BioRad #162-0177) or similar. Nitrocellulose membrane may also be used.
ECL Reagents	SuperSignal West Pico Chemiluminescent Substrate (Pierce #34080) or similar
NeutrAvidin-HRP	"High Sensitivity" (Pierce #31030).
Immunoprecipitation Kit (optional)	(Pierce #26146) or equivalent. Alternatively, commonly used IP reagents may be used.
Epoxomicin (optional)	(Boston Biochem # I-110) Cell-permeable proteasome inhibitor.

Sample Preparation

It is not in the scope of this protocol to guide the investigator on how and when a protein of interest may be ubiquitinated. It is assumed that a set of experimental conditions (including how many cells to use) has been defined by the investigator in which the potential ubiquitination of a protein will be examined.

1. Prepare cultured cells of choice by seeding either 6-well plates or tissue culture flasks as appropriate. Continue growth of cells in appropriate medium/conditions as previously established.
2. (Optional) At 4-12 hours prior to harvesting cells, add epoxomicin (or other cell-permeable proteasome inhibitor) to growth media at a final concentration of 5 μ M.
3. At 45 minutes prior to harvesting cells, thaw the 10X Ubl Lyse Reagent and the 100X DUB-Stop Protease Inhibitor Mix by gently and continuously swirling tubes in a lukewarm water bath ($\leq 30^{\circ}\text{C}$). Alternatively, tubes may be thawed with a rapid back-and-forth rolling motion between palms of hands. Do not heat tubes for an extended period of time. Once completely thawed, briefly centrifuge to collect contents at bottom of tubes. Do not store 100X DUB-Stop Mix on ice—it contains DMSO and will solidify if stored below room temperature. Wear gloves to prevent accidental exposure to DMSO.
4. For each cell lysis that will be done, prepare 1 ml of Lysis Buffer by combining the following in order (mix after each addition):
 - i. 890 μ l dH₂O
 - ii. 100 μ l 10X Ubl Lyse Reagent
 - iii. 10 μ l 100X DUB-Stop Mix

We recommend using 1 ml of Lysis Buffer per well of a 6-well tissue culture plate. Scale volumes appropriately for the number of lysates required and total amount of cells harvested. If more than 1 ml of Lysis Buffer is required, a 15 ml flip-top conical tube (such as a Falcon 2059) is convenient for mixing components. Vortex gently to mix components (avoid frothing), then chill the Lysis Buffer on ice for at least 20 minutes.

5. Harvest cells by aspirating culture media and rinsing once or twice with PBS. Cells may then be collected by standard trypsinization procedures to generate small cell pellets in centrifuge tubes. We recommend omitting the trypsinization process by directly adding 1 ml of the ice-cold Lysis Buffer to the rinsed, aspirated cells on the plates or flasks.

6. Use gentle, mechanical agitation to aid disruption of cells with the Lysis Buffer. Tapping the plates/flasks or repeatedly pipetting the Lysis Buffer over the surface of the cells helps to thoroughly solubilize the samples. The sample may become viscous due to release of genomic DNA. We recommend the use of 1 ml wide-bore pipette tips (made by cutting off the ends of the pipette tips with a sharp razor blade) to facilitate handling of the samples. Carefully transfer the samples to 1.5 ml microcentrifuge tube.
7. (Optional) Viscous lysates are not optimal for IP techniques. Genomic DNA may be sheered by a number of techniques in order to lower sample viscosity. We suggest using one of the following:
 - i. Sonication using a microtip. Care must be taken to avoid sample heating. Use multiple 10-15 second treatments at 50% duty, with icing for 1-2 minutes in between treatments to cool sample(s). 4-6 total cycles is effective in reducing viscosity. Be certain to rinse microtip between samples to avoid cross-contamination.
 - ii. Pass sample through a Qiashredder (Qiagen #79656) one or more times to reduce viscosity—see manufacturer's recommendations and protocols.
8. Clear insoluble material from samples by centrifugation at 4°C for 15-20 minutes at $\geq 15000 \times g$. Transfer supernatants to clean microfuge tubes and place on ice.
9. Samples are now ready for IP.

Immunoprecipitation of Target Protein

It is left to the investigator to determine how best to perform IP reactions. We recommend using a polyclonal antibody generated using a recombinant form of the target protein of interest. While monoclonal antibodies can also be used in the IP, the potential exists that the epitope recognized by the antibody may be a site of ubiquitination on the target protein. If so, then ubiquitinated target protein may not be IP'd, leading to a false-negative result. The efficacy of the antibody in the IP of the target protein should be previously validated by the investigator.

General suggested guidelines for target protein IP are as follows:

1. Transfer 400-1000 μ l of cell lysate(s) (obtained in Sample Preparation) to labeled, chilled 1.5 ml microcentrifuge tubes. Adjust the volume of lysate(s) used depending on the abundance of the target protein as previously determined by the investigator.
2. Add an appropriate amount of antibody to the sample and incubate for 1-4 hours at 4°C. We routinely begin with 20 μ g of polyclonal antibody.
3. Add 10-20 μ l (packed volume) of Protein A/G Sepharose ("resin") to the sample and place on rocker for 1-2 hours at 4°C (overnight is acceptable). For convenience, resin may be added as 20-40 μ l of a 50% slurry.
4. Place tubes containing 5X HS Wash Buffer and 10X LS Wash Buffer in a water bath ($\leq 30^{\circ}\text{C}$) or incubator until completely thawed. Gently vortex tubes to ensure solutions are well mixed. The 5X HS Buffer may be cloudy at room temperature.
5. For each IP reaction, make 2 ml of HS Wash Buffer by diluting 400 μ l of 5X stock into 1.6 ml of dH₂O. Make 1 ml of LS Wash Buffer by diluting 100 μ l of 10X stock into 0.9 ml of dH₂O. Scale as needed to generate enough of each wash buffer for all samples.
6. Chill both Wash Buffers on ice for at least 20 minutes.
7. Pellet resins from Step 3 by brief, low-speed centrifugation (1-2 minutes at 5000 xg). Carefully remove supernatant(s) by aspiration or pipetting.
8. Wash resin(s) by adding 1 ml ice-cold 1X HS Wash Buffer. Briefly vortex to suspend resin, then place tubes on a rocker or end-over-end mixer for 5 minutes at 4°C. Briefly centrifuge (1-2 minutes at 1000 xg) to pellet resin, then carefully aspirate wash supernatant. Repeat once for a total of two washes in 1X HS Wash Buffer.
9. Wash resin(s) twice with 0.5 ml ice-cold 1X LS Wash Buffer using the procedures listed in Step 8. After the last wash, carefully remove all supernatant by aspiration.
10. Add 30 μ l of 2X SDS-PAGE sample buffer and 4 μ l 1M DTT. Mix well and heat reactions to 80-90°C for 5 minutes. Centrifuge briefly to pellet resin.

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565

www.bostonbiochem.com

SDS-PAGE and Western Blot Analysis

The choice of SDS-PAGE gel that is used for separating the target protein of interest will depend on the size of the target protein. A constant percentage or gradient gel may be used. We routinely use BioRad's *Criterion* gels and buffering systems which are available in many formats and yield consistent Western Blot results.

General suggested guidelines for SDS-PAGE/Western Blot of target protein IP are as follows:

1. Assemble SDS-PAGE gel according to manufacturer's instructions.
2. Thaw the Ubiquitinated Positive Control sample included with the kit. Add 4 μ l of 1M DTT, mix, and heat at 37-40 °C for 10 minutes. (Enough Positive Control is provided for approximately 4 lanes worth of analysis. Only add DTT to the Positive Control the first time it is opened. Freeze down any unused Positive Control sample at -80°C in useful aliquot sizes for future use.)
3. Load 10 μ l of Positive Control into the appropriate gel well(s).
4. Load 10-30 μ l of each sample (from Step 10 in Immunoprecipitation of Target Protein section) into appropriate wells. The volume loaded will depend on your gel format.
5. Run the gel until the dye-front just reaches bottom of gel (approximately 1 hour at 180V using the *Criterion* gel system—adjust run times and voltage accordingly for your system).
6. Carefully disassemble gel and transfer to a small tray containing 50-100 ml of Towbin Buffer. Soak the gel at room temperature for 20 minutes with gentle rocking. Some gels tend to float on top of the buffer—this should be minimized with gentle agitation.
7. Prepare PVDF membrane by wetting in a small amount of 100% methanol for 1-2 minutes. When membrane is completely wetted, transfer to 20-30 ml of Towbin Buffer for 5 minutes at room temperature with gentle rocking.
8. Assemble transfer "sandwich" using pre-wetted blotting paper, gel, PVDF membrane, and any other components for electroblotting according to system manufacturer's suggested protocol. (We utilize the BioRad Trans•Blot SD Semi Dry Transfer Cell)
9. Electroblot gel contents to PVDF at 15V (constant) for 45 minutes at room temperature (settings will depend on your system).
10. Disassemble gel sandwich and prepare for Blocking/Antibody Detection.

11. Prepare Blocking Solution by dissolving 3 grams solid Fraction V BSA in 100 ml PBST solution (PBST+3% w/v BSA). The Blocking Solution may be filter sterilized and stored at 4°C for up to 5 days, but we suggest preparing it fresh when needed.
12. Soak PVDF membrane in 50-100 ml Blocking Solution 1 hour at room temperature (or overnight at 4°C) with gentle rocking in a covered container or sealable bag.
13. Decant Blocking Solution and rinse membrane for 10-15 minutes in 50 ml PBST.
14. Prepare antibody solution by diluting 40 µl of biotinylated α -Ubiquitin antibody (supplied) in 4 ml of PBST + 0.5% w/v BSA.
15. Using a sealable plastic bag, incubate membrane in antibody solution for 2 hours at room temperature (or overnight at 4°C) with rocking or shaking. Make certain that the entire PVDF membrane surface is uniformly exposed to antibody solution.
16. Remove membrane from antibody solution and wash with 50-100 ml PBST for 15 minutes at room temperature using rocking or shaking. Decant wash solution. Repeat twice for a total of three membrane washes.
17. Prepare avidin-HRP solution by diluting 4 µl of Pierce High Sensitivity NeutrAvidin HRP (or equivalent) in 20 ml of PBST + 0.5% BSA.
18. Incubate membrane in avidin-HRP solution for 60 minutes at room temperature (longer incubations may lead to higher background signals) with rocking or shaking. Make certain that the entire PVDF membrane surface is uniformly exposed to solution.
19. Remove membrane from avidin-HRP solution and wash with 50-100 ml PBST for 15 minutes at room temperature using rocking or shaking. Decant wash solution. Repeat twice for a total of three washes.
20. Membrane is now ready for ECL detection procedure. Chemiluminescence reagents for detecting products are available from a number of sources. We recommend using SuperSignal West Pico Chemiluminescent Substrate from Pierce. Typical film exposure times range from 5 seconds to 2 minutes. Using the protocol listed above, the Positive Control is typically observed with film exposures of 60 seconds or less.

Example Data




Figure 1. Ubi-Detect Analysis of Polyubiquitinated HDM2

In vitro reactions using purified ubiquitin ligase GST-HDM2, ubiquitin, Mg^{2+} -ATP and cellular fractions from HCT-116 cells were used to generate autoubiquitinated starting material for analysis. Reactions were terminated with an excess volume of Ubl-Lyse reagent containing DUB-Stop Protease Inhibitor Mix. Using methods described in this protocol, protein was immunoprecipitated using either an anti-HDM2 antibody (R&D Systems # MAB1244) or a negative control antibody ("Con"). IP'd complexes were resolved on 4-20% SDS-PAGE gels and then electroblotted to PVDF membranes. Left Panel: membrane was developed using anti-MDM2 rabbit pAb (R&D Systems # AF1244) at 0.05 mg/ml and anti-rabbit HRP (R&D Systems HAF 008) at 1/2000 dilution. IP'd GST-HDM2 is detected as a discrete band (unmodified) and a high molecular weight smear. Right Panel: membrane was developed as described in the protocol. The presence of ubiquitinated GST-HDM2 is confirmed.

Literature

- References:** Choo, Y.S. & Zhang, Z. (2009) *J. Vis Exp.* **30**: 1293
 Laney, J.D. & Hochstrasser, M. (2011) *Curr. Proto. Prot. Sci.*
 DOI:10.1002/0471140864.ps1405s29

For Laboratory Research Use Only, Not For Use in Humans

Rev: 05/05/2014

840 Memorial Drive, Cambridge, MA 02139 Phone: 617-576-2210 FAX: 617-492-3565
www.bostonbiochem.com

The contents of this datasheet (unless otherwise noted) are Copyright © 2008 Boston Biochem, Inc. All rights reserved. Duplication in whole or in part is strictly prohibited without the express written consent of Boston Biochem, Inc. "Boston Biochem" is a Trademark of Boston Biochem, Inc., 840 Memorial Drive, Cambridge, MA 02139.