

Human Siglec-3/CD33 LlamabodyTM V_HH His-tag Alexa Fluor® 647-conjugated Antibody

Recombinant Monoclonal Llama V_HH domain Clone # L008.2.14N

Ostalas Nusskas I EAD40700D

| DESCRIPTION | | | |
|--------------------|---|--|--|
| Species Reactivity | Human | | |
| Specificity | Detects human Siglec-3/CD33 in direct ELISAs. | | |
| Source | Recombinant Monoclonal Llama V _H H domain Clone # L008.2.14N | | |
| Purification | His-tag purified from cell culture supernatant | | |
| Immunogen | Chinese Hamster Ovary cell line CHO-derived human Siglec-3/CD33 Val18-His259 Accession # AAA51948 | | |
| Conjugate | Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm | | |
| Formulation | Supplied in a saline solution containing BSA and Sodium Azide. | | |

*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

| APPLICATIONS | | |
|---|---------------------------------------|--|
| Please Note: Optimal dilutions should be determined | by each laboratory for each applicati | on. General Protocols are available in the Technical Information section on our website. |
| | Recommended Concentration | Sample |
| Flow Cytometry | 0.25-1 μg/10 ⁶ cells | HEK293 human embryonic kidney cell line transfected with human Siglec-3/CD33 and eGFP |

| DATA | | | |
|---------------------|---|--|--|
| Flow Cytometry | Detection of Siglec-3/CD33 in HEK293 Human Cell Line Transfected with Human Siglec-3/CD33 and eGFP by Flow Cytometry. HEK293 human embryonic kidney cell line transfected with human Siglec- 3/CD33 and eGFP was stained with (A) Llama Anti-Human Siglec-3/CD33 Alexa Fluor® 647- conjugated Llamabody VHH Monoclonal Antibody (Catalog # LFAB10733R) or normal Ilama IgG (Catalog # IC011R). Staining was performed using our Staining Membrane-associated Proteins protocol. | | |
| PREPARATION AND S | STORAGE | | |
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. | | |
| Stability & Storage | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied. | | |

Rev. 7/27/2022 Page 1 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449



Human Siglec-3/CD33 LlamabodyTM V_HH His-tag Alexa Fluor® 647-conjugated Antibody

Recombinant Monoclonal Llama V_HH domain Clone # L008.2.14N

BACKGROUND

Siglecs (sialic acid binding Ig-like lectins) are I-type (Ig-type) lectins belonging to the Ig superfamily. They are characterized by an N-terminal Ig-like V-type domain which mediates sialic acid binding, followed by varying numbers of Ig-like C2-type domains (1, 2). Eleven human Siglecs have been cloned and characterized. They are sialoadhesin/CD169/Siglec-1, CD22/Siglec-2, CD33/Siglec-3, Myelin-Associated Glycoprotein (MAG/Siglec-4a) and Siglecs 5 to 11 (1-3). To date, no Siglec has been shown to recognized any cell surface ligand other than sialic acids, suggesting that interactions with glycans containing this carbohydrate are important in mediating the biological functions of Siglecs. Siglecs 5 to 11 share a high degree of sequence similarity with CD33/Siglec-3 both in their extracellular and intracellular regions. They are collectively referred to as CD33-related Siglecs. One remarkable feature of the CD33-related Siglecs is their differential expression pattern within the hematopoietic system (1, 2). This fact, together with the presence of two conserved immunoreceptor tyrosine-based inhibition motifs (ITIMs) in their cytoplasma tails, suggests that CD33related Siglecs are involved in the regulation of cellular activation within the immune system. Human Siglec-3 is alternatively known as myeloid cell surface antigen CD33 and GP67. Human Siglec-3 cDNA encodes a 364 amino acid (aa) polypeptide with a hydrophobic signal peptide, an N-terminal Ig-like V-type domain, one Ig-like C2-type domains, a transmembrane region and a cytoplasmic tail (1, 4). Siglec-3 expression is restricted to cells of myelomonocytic lineage (2). It binds sialic acid preferring a2,3-linkage over a2,6-linkage (5). Studies indicated that Siglec-3 can mediate inhibitory signals (7).

References:

- 1. Crocker, P.R. and A. Varki (2001) Trends Immunol. 22:337.
- 2. Crocker, P.R. and A. Varki (2001) Immunology 103:137.
- 3. Angata, T. et al. (2002) J. Biol. Chem. 277:24466.
- 4. Simmons, D. and B. Seed (1988) J. Immunol. 141:2797.
- 5. Freeman, S.D. et al. (1995) Blood 85:2002.
- 6. Taylor, V.C. et al. (1999) J. Biol. Chem. 274:11505.
- 7. Ulyanova, T. et al. (1999) Eur. J. Immunol. 29:3440

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 7/27/2022 Page 2 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449