

DESCRIPTION


Species Reactivity	Human
Specificity	Recognizes human Leukotriene B4 R1 (Pettersson, A. <i>et al.</i> (2000) <i>Biochem. Biophys. Res. Commun.</i> 279 :520).
Source	Monoclonal Mouse IgG ₁ Clone # 203/14F11
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	HeLa cervical epithelial carcinoma cell line transfected with human Leukotriene B4 R1
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Polymorphonuclear granulocytes secrete the lipid chemotactic mediator Leukotriene B4 (LTB₄) in response to inflammatory stimuli (1). Neutrophils, monocytes and lymphocytes respond to LTB₄ via specific receptors localized on the cell surface (2-4). The high affinity LTB₄ Receptor known as BLT1 is only expressed on leukocytes (5-7) while a second low affinity receptor BLT2 is expressed more ubiquitously (8, 9). The BLT1 and BLT2 are G-protein linked seven-transmembrane spanning receptors that share about 37-45% amino acid identity (8, 9).

References:

1. Samuelsson, B. *et al.* (1987) *Science* **237**:1171.
2. Brom, J. and W. Konig (1989) *Immunology* **68**:479.
3. Patry, C. *et al.* (1996) *Prostaglandins, Leukotrienes and Essential Fatty Acids* **54**:361.
4. Dasari, V.R. *et al.* (2000) *Immunopharm.* **48**:157.
5. Pettersson, A. *et al.* (2000) *Biochem. Biophys. Res. Comm.* **279**:520.
6. Yokomizo, T. *et al.* (1997) *Nature* **387**:620.
7. Kato, K. *et al.* (2000) *J. Exp. Med.* **192**:413.
8. Yokomizo, T. *et al.* (2000) *J. Exp. Med.* **192**:421.
9. Kamohara, M. *et al.* (2000) *J. Biol. Chem.* **275**:27000.