

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human VSTM-2A in direct ELISAs.
Source	Monoclonal Mouse IgG _{2B} Clone # 1024718
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Human embryonic kidney cell HEK293-derived human VSTM2A Ser25-Phe244 Accession # Q8TAG5-2
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA	This antibody functions as an ELISA capture antibody when paired with Mouse Anti-Human VSTM2A Monoclonal Antibody (Catalog # MAB100371).
	This product is intended for assay development on various assay platforms requiring antibody pairs.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

V-set and transmembrane domain-containing protein 2A (VSTM2A) is a secreted glycoprotein that is expressed by committed preadipocytes. N-linked glycosylation is crucial for its secretion, but not for preadipocyte cell differentiation activity. It is expressed during adipocyte development and its over-expression promotes adipogenesis (1). VSTM2A is highly expressed in the brain and *Vstm2a* was identified as an enigmatic gene that is highly produced in mouse brain (1, 2). A positive association has been observed between *Vstm2a* and *Pparg2*. VSTM2A plays a role in the regulation of the early stage of white and brown preadipocyte cell differentiation. It promotes adipogenic commitment of preadipocytes by increasing gene expression of the transcription factor PPARG in a BMP4-dependent signaling pathway (1, 3). In humans, two isoforms (1 and 2) exist due to alternative splicing. Human VSTM2A is synthesized either as a 236 amino acid (aa) (Isoform 1) or a 244 aa (Isoform 2) precursor that contains a 24 aa signal sequence followed by the VSTM2A domain. Human Isoform 2 VSTM2A shares 78% aa sequence identity with mouse and rat VSTM2A.

References:

1. Secco, B. *et al.* (2017) Cell Rep. **18**:93.
2. Pandey, A.K. *et al.* (2014) PloS One. **9**:e88889.
3. Berry, D.C. *et al.* (2013) Development. **140**:3939.