

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human ADAM8 in direct ELISAs and Western blots. In Western blots, less than 5% cross-reactivity with the ectodomains of recombinant human ADAM9, 10, 15, 17, and 19 is observed. Detects an epitope within aa 201-497 of recombinant human ADAM8 which corresponds to the metalloprotease and disintegrin-like domains.
<b>Source</b>	Monoclonal Mouse IgG <sub>2B</sub> Clone # 143303
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human ADAM8 ectodomain Asp158-Ser653 Accession # P78325
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
<b>Western Blot</b>	1 µg/mL	See Below
<b>Immunoprecipitation</b>	25 µg/mL	Conditioned cell culture medium spiked with Recombinant Human ADAM8 aa 158-497 (Catalog # 1031-AD), <a href="#">see our available Western blot detection antibodies</a>

## DATA



## PREPARATION AND STORAGE

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

ADAM8, also known as cell surface antigen MS2 or CD156a, is a member of the ADAM family that contains a disintegrin and metalloprotease-like domain (1, 2). ADAM8 can cleave a variety of substrates and has been shown as a sheddase for the low affinity IgE receptor CD23 and the neural recognition molecule CHL1 (3, 4). Expression and regulation studies suggest that ADAM8 is a novel osteoclast stimulating factor and may play a role in asthma (5, 6). The 824 amino acid precursor of human ADAM8 consists of a signal peptide (residues 1 to 16), a pro peptide (residues 17 to 199), a metalloprotease domain (residues 200 to 400), a disintegrin-like domain (residues 408 to 494), a cysteine-rich region (residues 497 to 613), an EGF-like domain (residues 614 to 640), a transmembrane region (residues 656 to 676) and a cytoplasmic domain (residues 677 to 824).

**References:**

1. Yoshiyama, K. *et al.* (1997) *Genomics* **41**:56.
2. Moss, M.L. and J.W. Bartsch (2004) *Biochemistry* **43**:7227.
3. Fourie, A.M. *et al.* (2003) *J. Biol. Chem.* **278**:30469.
4. Naus, S. *et al.* (2004) *J. Biol. Chem.* **279**:16083.
5. Choi, S.J. *et al.* (2001) *J. Bone Miner. Res.* **16**:814.
6. King, N.E. *et al.* (2004) *Am. J. Respir. Cell Mol. Biol.* **31**:257.