

**DESCRIPTION**


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human Sonic Hedgehog/Shh in direct ELISAs.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 605022
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived human Sonic Hedgehog/Shh protein Cys24-Gly197 Accession # NP_000184.1
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

<b>ELISA</b>	<p>This antibody functions as an ELISA detection antibody when paired with Mouse Anti-Human Sonic Hedgehog/Shh Monoclonal Antibody (Catalog # <a href="#">MAB10939</a>).</p> <p>This product is intended for assay development on various assay platforms requiring antibody pairs. We recommend the Human Sonic Hedgehog/Shh N-Terminus DuoSet ELISA Kit (Catalog # <a href="#">DY1314-05</a>) for convenient development of a sandwich ELISA.</p>
--------------	---

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.5 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<p><b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b></p> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

Sonic Hedgehog (Shh) is expressed in embryonic tissues that are critical for the patterning of the developing central nervous system, somite, and limb. It is also involved in whisker, hair, foregut, tooth, and bone development. Shh regulates neural and hematopoietic stem cell fate and is important for thymocyte differentiation and proliferation as well as T cell determination. In adult tissue Shh is associated with cancer development and tissue remodeling following injury (1-3). Human Shh encodes a 462 amino acid (aa) precursor protein that is autocatalytically processed to yield a non-glycosylated 19 kDa N-terminal fragment (Shh-N) and a glycosylated 25 kDa C-terminal protein (Shh-C) (4). Shh-C, which is responsible for the intramolecular processing of Shh, is rapidly degraded following Shh proteolysis (5). Shh-N is highly conserved, sharing >98% aa identity between mouse, human, rat, canine, porcine, and chicken Shh-N. Shh-N can be palmitoylated at its N-terminal cysteine and modified by cholesterol addition at its C-terminus (6). These modifications contribute to the membrane tethering of Shh as well as its assembly into various sized multimers (6-9). Lipid modification and multimerization greatly increase Shh-N receptor binding affinity and signaling potency (5, 6, 8, 9). Monomeric and multimeric Shh can be released from the plasma membrane by the cooperative action of DISP1, SCUBE2, and TACE/ADAM17 (10-12). Modifications also extend the effective range of Shh functionality and are required for the development of protein gradients important in tissue morphogenesis (9, 13). Canonical signaling of Shh is mediated by a multicomponent receptor complex that includes Patched (PTCH1, PTCH2) and Smoothened (SMO) (14). The binding of Shh to PTCH releases the basal repression of SMO by PTCH. Shh activity can also be regulated through interactions with heparin, glypicans, and membrane-associated Hip (hedgehog interacting protein) (13, 15, 16).

**References:**

1. Briscoe, J. and P.P. Thérond (2013) *Mol. Cell. Biol.* **14**:416.
2. Aviles, E.C. *et al.* (2013) *Front. Cell. Neurosci.* **7**:86.
3. Xie, J. *et al.* (2013) *OncoTargets Ther.* **6**:1425.
4. Marigo, V. *et al.* (1995) *Genomics* **28**:44.
5. Zeng, X. *et al.* (2001) *Nature* **411**:716.
6. Feng, J. *et al.* (2004) *Development* **131**:4357.
7. Goetz, J.A. *et al.* (2006) *J. Biol. Chem.* **281**:4087.
8. Pepinsky, R.B. *et al.* (1998) *J. Biol. Chem.* **273**:14037.
9. Chen, M.-H. *et al.* (2004) *Genes Dev.* **18**:641.
10. Etheridge, L.A. *et al.* (2010) *Development* **137**:133.
11. Jakobs, P. *et al.* (2014) *J. Cell Sci.* **127**:1726.
12. Dierker, T. *et al.* (2009) *J. Biol. Chem.* **284**:8013.
13. Lewis, P.M. *et al.* (2001) *Cell* **105**:599.
14. Carpenter, D. *et al.* (1998) *Proc. Natl. Acad. Sci. USA* **95**:13630.
15. Filmus, J. and M. Capurro (2014) *Matrix Biol.* **35**:248.
16. Chuang, P.-T. and A.P. McMahon (1999) *Nature* **397**:617.