

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Hepassocin/FGL1 in direct ELISA.
Source	Monoclonal Mouse IgG ₁ Clone # 1052924
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Hepassocin/FGL1 Met1-Ile312 Accession # BAB70690
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

ELISA	This antibody functions as an ELISA detection antibody when paired with mouse anti-Hepassocin/FGL1 Monoclonal Antibody (Catalog # MAB11230). This product is intended for assay development on various assay platforms requiring antibody pairs.
--------------	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Hepassocin, also known as hepatocyte-derived fibrinogen-related protein 1 (HFREP-1) and FGL1, is a liver-specific secreted protein belonging to the fibronogen superfamily, whose members share a fibrinogen domain at their C-termini. Hepassocin may play a role in the development of hepatocellular carcinomas.