Human HLA-DR Antibody

Recombinant Monoclonal Rabbit IgG Clone # 3359B
Catalog Number: MAB11834

DESCRIPTION

Species Reactivity	Human
Specificity	Detects a synthetic peptide specific for human HLA-DR around amino acid 215 in Direct ELISA.
Source	Recombinant Monoclonal Rabbit IgG Clone # 3359B
Purification	Protein A or G purified from cell culture supernatant
Immunogen	Synthetic Peptide Accession # P01903
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose.

APPLICATIONS

Please Note: *Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.*

	Recommended Concentration	Sample
Western Blot	0.5 µg/mL	HDLM-2 human Hodgkin's lymphoma cell line
Immunohistochemistry	0.25-25 µg/mL	Immersion fixed paraffin-embedded sections of tonsil
COMET	0.5 µg/mL	Immersion fixed paraffin-embedded sections of human tonsil


Bio-Techne®
Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956
USA | TEL: 800.343.7475 Canada | TEL: 855.668.8722 Europe | Middle East | Africa TEL: +44.0.1235.529449
China | info.cn@bio-techne.com TEL: 400.821.3475

Human HLA-DR Antibody

Recombinant Monoclonal Rabbit IgG Clone # 3359B
Catalog Number: MAB11834


DATA

Multiplex Immunofluorescence




HLA-DRA in Human Tonsil via seqIF™ staining on COMET™
HLA-DRA was detected in immersion fixed paraffin-embedded sections of human tonsil using Rabbit Anti-Human HLA-DRA, Monoclonal Antibody (Catalog # MAB11834) at 0.5ug/mL at 37 ° Celsius for 4 minutes. Before incubation with the primary antibody, tissue underwent an all-in-one dewaxing and antigen retrieval preprocessing using PreTreatment Module (PT Module) and Dewax and HIER Buffer H (pH 9; Epredia Catalog # TA-999-DHBH). Tissue was stained using the Alexa Fluor™ Plus 647 Goat anti-Rabbit IgG Secondary Antibody at 1:200 at 37 ° Celsius for 2 minutes. (Yellow; Lunaphore Catalog # DR647RB) and counterstained with DAPI (blue; Lunaphore Catalog # DR100). Specific staining was localized to the membrane. Protocol available in COMET™ Panel Builder.

Immunohistochemistry



Immersion fixed paraffin-embedded sections of tonsil
HLA-DR was detected in immersion fixed paraffin-embedded sections of human tonsil using Rabbit Anti-Human HLA-DR Monoclonal Antibody (Catalog # MAB11834) at 1 µg/ml overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using VisUCyte Antigen Retrieval Reagent-Basic (Catalog # VCTS021). Tissue was stained using the HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # HAF008) and counterstained with hematoxylin (blue). Specific staining was localized to the membrane. View our protocol for Chromogenic IHC Staining of Paraffin-embedded Tissue Sections.

Western Blot



Detection of Human HLA-DR by Western Blot. Western Blot shows lysates of HDLM-2 human Hodgkin's lymphoma cell line. PVDF membrane was probed with 0.5 µg/ml of Rabbit Anti-Human HLA-DR Monoclonal Antibody (Catalog # MAB11834) followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # HAF008). A specific band was detected for HLA-DR at approximately 35 kDa (as indicated). This experiment was conducted under reducing conditions and using Western Blot Buffer Group 1.

PREPARATION AND STORAGE

Reconstitution	Reconstitute lyophilized material at 0.2 mg/ml in sterile PBS. For liquid material, refer to CoA for concentration.
Shipping	Lyophilized product is shipped at ambient temperature. Liquid small pack size (-SP) is shipped with polar packs. Upon receipt, store immediately at the temperature recommended below.
Stability & Storage	**Use a manual defrost freezer and avoid repeated freeze-thaw cycles.** • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

HLA-DR is a transmembrane human major histocompatibility complex 2 (MHC II) family member and consists of a 34 kDa (alpha) subunit and one of several 28 kDa (beta) subunits. HLA-DR is expressed primarily by B cells and dendritic cells (DC), in which it binds peptides derived from internalized and processed antigenic proteins. It presents these peptides on the cell surface for recognition by the T cell receptor on $CD4^+$ T cells. This interaction is central to antigen specificity in the adaptive immune response. HLA-DR alleles, polymorphisms, and aberrant expression are linked to a variety of diseases including autoimmunity and cancer.

Bio-Techne®
Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956
USA | TEL: 800.343.7475 Canada | TEL: 855.668.8722 Europe | Middle East | Africa TEL: +44.0.1235.529449
China | info.cn@bio-techne.com TEL: 400.821.3475