RD SYSTEMS a biotechne brand

Recombinant Monoclonal Rabbit IgG Clone # 2622G Catalog Number: MAB12490

 DESCRIPTION

 Species Reactivity
 Human

 Specificity
 Detects human GLP-1 in direct ELISAs.

 Source
 Recombinant Monoclonal Rabbit IgG Clone # 2622G

 Purification
 Protein A or G purified from cell culture supernatant

 Immunogen
 Human GLP-1 synthetic peptide Accession # P01275

 Formulation
 Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

 Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

 ELISA
 This antibody functions as an ELISA detection antibody when paired with Rabbit Anti-Human GLP-1 Monoclonal Antibody (Catalog # MAB124901).

PREPARATION AND STORAGE	
Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	 Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Glucagon is a preprotein which is cleaved into four distinct peptides, including the hormone GLP-1 (aa 98-128). GLP-1 is a secreted hormone with multiple effects upon the intestine (gastric motility), pancreas (glucose dependent insulin release) and hypothalamic pituitary axis (modulates LH, THS, CRH, ocytocin and vasopressin secretion). It also affects plasma glucagon levels. GLP-1 is expressed in enteroendocrine L cells and neurons of the caudal brainstem which project to the forbain, the amygdala and the hypothalamus. Recent studies show pancreatic intra-islet GLP-1 expression, which is regulated by cytokines, hyperglycemia and cell injury.

Rev. 1/31/2020 Page 1 of 1



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 **Canada** TEL 855 668 8722 **China** TEL +86 (21) 52380373 **Europe | Middle East | Africa** TEL +44 (0)1235 529449