

DESCRIPTION



Species Reactivity	Human
Specificity	Detects human α -Synuclein in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 970114
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human α -Synuclein Met1-Ala140 Accession # P37840
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	5 μ g/mL	See Below
Immunohistochemistry	5-25 μ g/mL	See Below

DATA

<p>Western Blot</p>  <p>Detection of Human α-Synuclein by Western Blot. Western blot shows lysates of human brain (motor cortex and cerebellum) tissue. PVDF membrane was probed with 5 μg/mL of Mouse Anti-Human α-Synuclein Monoclonal Antibody (Catalog # MAB13383) followed by HRP-conjugated Anti-Mouse IgG Secondary Antibody (Catalog # HAF018). A specific band was detected for α-Synuclein at approximately 17 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 3.</p>	<p>Immunohistochemistry</p>  <p>α-Synuclein in Human Brain. α-Synuclein was detected in immersion fixed paraffin-embedded sections of human brain (hippocampus) using Mouse Anti-Human α-Synuclein Monoclonal Antibody (Catalog # MAB13383) at 5 μg/mL for 1 hour at room temperature followed by incubation with the Anti-Mouse IgG VisUCyte™ HRP Polymer Antibody (Catalog # VC001). Tissue was stained using DAB (brown) and counterstained with hematoxylin (blue). Specific staining was localized to synaptic boutons. View our protocol for IHC Staining with VisUCyte HRP Polymer Detection Reagents.</p>
--	---

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.5 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

α -Synuclein is a cytoplasmic protein that is predominantly expressed in the central nervous system. It is a precursor of a minor peptide found in amyloid deposition that is a hallmark of Alzheimer's disease. Defects in α -Synuclein are associated with familial Parkinson's disease.