

Human Lymphotoxin Antibody

Monoclonal Mouse IgG₁ Clone # 135125 Catalog Number: MAB1370

DESCRIPTION	
Species Reactivity	Human
Specificity	Recognizes human Lymphotoxin α chains in α ₂ β ₁ or α ₁ β ₂ trimers and also human TNF-β. In direct ELISAs, less than 10% cross-reactivity with recombinant human (rh) APRIL, rhBAFF, rhEDA-A2, rhFas Ligand, rhLIGHT, rhGITR Ligand, rhOX40 Ligand, recombinant mouse RANKL, rhTNF-α, rhTRAIL, rhTRANCE, rhTWEAK, or rhVEGI is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 135125
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	S. frugiperda insect ovarian cell line Sf 21-derived recombinant human Lymphotoxin α/β
Endotoxin Level	<0.10 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.		
Neutralization	Measured by its ability to neutralize Lymphotoxin α2/β1induced cytotoxicity in the L929 mouse fibroblast cell line. The Neutralization Dose (ND50) is typically 0.150 - 3.00 μg/mL in the presence of 10 ng/mL Recombinant Human	
	Lymphotoxin α2/β1and actinomycin D.	



- Use a manual denost neezer and avoid repeated neeze-thaw cycle
 - 12 months from date of receipt, -20 to -70 °C as supplied.
 1 month. 2 to 8 °C under sterile conditions after reconstitution.
 - I month, 2 to 8 C under sterile conditions after reconstitution.
 C months, 20 to 70 °C under sterile conditions ofter reconstitution.
 - 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Lymphotoxin α , also known as tumor necrosis factor (TNF) β , and Lymphotoxin β are members of the TNF superfamily and have been designated TNFSF1 and TNFSF3, respectively. Secreted Lymphotoxin α forms a homotrimer (TNFbeta) and heterotrimeric complexes with the type II membrane protein, Lymphotoxin β (either $\alpha 2\beta 1$ or $\alpha 1\beta 2$ timers). These trimers serve as ligands for Lymphotoxin β receptor, TNF RI and TNF RII and play a role in the development and maturation of the immune system.

References:

- 1. Matthews, N. and M. L. Neale (1987) in Lymphokines and Interferons, A Practical Approach.
- 2. Clemens, M. J. et al. (eds): IRL Press. 221.

Rev. 7/18/2024 Page 1 of 1

Bio-Techne®

Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956 USA | TEL: 800.343.7475 Canada | TEL: 855.668.8722 Europe | Middle East | Africa TEL: +44.0.1235.529449 China | info.cn@bio-techne.com TEL: 400.821.3475